

Generalized logarithmic derivatives for K_n

Sarah Livia Zerbes

February 2, 2008

dedicated to John Coates

Abstract

We construct (generalized) logarithmic derivatives for general n -dimensional local fields $\mathcal{K}$ of mixed characteristics $(0, p)$ in which p is not necessarily a prime element with residue field k such that $[k : k^p] = p^{n-1}$. For the construction of the logarithmic derivative map, we define n -dimensional rings of overconvergent series and show that - as in the 1-dimensional case - they can be interpreted as functions converging on some annulus of the open unit p -adic disc. Using the generalized logarithmic derivative, we give a new construction of Kato's n -dimensional dual exponential map.

2000 *Mathematics Subject Classification*. Primary 11S70; Secondary 19F99 11R23

Contents

1	Introduction	2
1.1	The main results	2
1.2	Acknowledgements	3
2	Setup	3
2.1	Fields of norms	3
2.2	Lift to characteristic 0	5
2.3	Some details about $\phi(\pi_K)$	6
3	Some stuff we need	10
3.1	Milnor K -groups and completions	10
3.2	Modules of differentials	10
3.3	The norm map	12
4	The logarithmic derivative	14
4.1	Some more modules of differentials	14
4.2	The ψ -operator	15
4.3	The logarithmic derivative	16

5	Higher dimensional rings of overconvergent series	18
5.1	Definition and properties	18
5.1.1	Definition of $\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}$	18
5.1.2	Definition of $\mathcal{R}_K^{\dagger, N}$	20
5.1.3	Properties of $\mathcal{R}_K^{\dagger}$	21
5.2	Properties of ϕ	23
5.3	Modules of differentials	24
6	The logarithmic derivative revisited	26
7	Relation to the dual exponential map	27
7.1	Statement of the theorem	27
7.2	Proof	28
7.2.1	Evaluation of the logarithmic derivative	28
7.2.2	The maps $f_{(\zeta_{p^i})_{i \geq 1}, m}$	30
7.2.3	Lots of commutative diagrams	31

1 Introduction

1.1 The main results

Let p be any prime number and F a finite unramified extension of $\mathbb{Q}_p$. For each $n \geq 1$, let μ_{p^n} be a primitive p^n th root of unity such that $\mu_{p^n}^p = \mu_{p^{n-1}}$, and let $F_n = F(\mu_{p^n})$. Let $F_\infty = F(\mu_{p^\infty})$ and $\Gamma = \text{Gal}(F^{\text{cyc}}/F)$, where $F^{\text{cyc}} \subset F_\infty$ is the cyclotomic $\mathbb{Z}_p$ -extension of F . One of the major tools in classical Iwasawa theory are so-called Coleman power series (which were discovered by Coates and Wiles [4] and Coleman [5]), which associate to every norm-compatible system of units $u = (u_n)$ in the tower $F_\infty = \bigcup_n F_n$ a norm-invariant element $G_u(T)$ in the Iwasawa algebra $\Lambda(\Gamma) \cong O_F[[T]]$ such that $G_u(\mu_{p^n}) = u_n$. It was unknown for a long time how to generalize this construction for finite *ramified* extensions of $\mathbb{Q}_p$, and only recently Cherbonnier and Colmez [3] discovered a way to construct Coleman series for such extensions using a formal analogy between $\Lambda(\Gamma)$ and Fontaine's rings of overconvergent series.

In [8], Fukaya extends the classical construction of Coleman power series to a certain class of 2-dimensional local fields of mixed characteristics $(0, p)$ with imperfect residue fields. Given such a field $\mathcal{H}$, she replaces the cyclotomic $\mathbb{Z}_p$ -extension by a certain 2-dimensional false Tate curve extension $\mathcal{H}_\infty$ of $\mathcal{H}$ (the field $\mathcal{H}_\infty$ is the direct limit of a sequence of finite Galois extensions $\mathcal{H}_n$ of $\mathcal{H}$), and in analogy to the classical case, her Coleman isomorphism associates to every element in $\varprojlim K_2(O_{\mathcal{H}_n})$ a norm-invariant element in $K_2(O_{\mathcal{H}}[[T]])$. Composing this map with the $d \log$ -operator, she obtains a canonical homomorphism

$$\tau : \varprojlim K_2(O_{\mathcal{H}_n}) \longrightarrow (\Omega_{O_{\mathcal{H}}[[T]]}^2)^{\psi=1}, \quad (1)$$

where ψ is the derivative of the norm map. Because of its analogy with the classical case, this map is called the 2-dimensional logarithmic derivative.

However, Fukaya needs to assume for her construction that p is a prime element in $\mathcal{H}$. Using the theory of higher-dimensional fields of norms developed by Scholl in [13], in this paper we define n -dimensional rings of overconvergent series, and following the approach of Cherbonnier and Colmez in [2], we use these rings to extend Fukaya's construction of the logarithmic derivative map to the case of a ramified basefield and to n -dimensional local fields of mixed characteristic $(0, p)$. The main result of this paper is the following theorem:-

Theorem 55. The logarithmic derivative gives a canonical homomorphism

$$\mathfrak{T} : \varprojlim \widehat{K}_n(\mathcal{K}_i) \longrightarrow (\Omega_{S_K^\dagger, N}^n(\log))^{\psi=1}. \quad (2)$$

In Section 7, we then use the logarithmic derivative map to give a new construction of Kato's dual exponential map for K_n .

1.2 Acknowledgements

I am very grateful to John Coates for his interest and to him and Yann Weil for their encouragement whenever I got fed up with the stubborn K -groups. Also, I would like to thank Max Karoubi for helpful discussions, Takako Fukaya for sending me copies of her paper [7] and Tony Scholl for giving me his preprint [13] on higher-dimensional local fields.

2 Setup

2.1 Fields of norms

Throughout this paper, we let p be some prime > 2 . (The calculations in the case $p = 2$ are slightly more difficult and will be dealt with in another paper.)

Let K be a finite totally ramified extension of $\mathbb{Q}_p$ with ring of integers O_K (we allow the case when $K = \mathbb{Q}_p$), and assume that $K \cap \mathbb{Q}_p(\mu_{p^\infty}) = \mathbb{Q}_p$. We construct our n -dimensional local field inductively:- Let

$$B_1 = O_K[[T_1]][T_1^{-1}]^\vee, \quad (3)$$

$$B_i = B_{i-1}[[T_i]][T_i^{-1}]^\vee \text{ for } 2 \leq i < n, \quad (4)$$

and define

$$O_{\mathcal{K}} = B_{n-1}, \quad (5)$$

$$\mathcal{K} = O_{\mathcal{K}}[p^{-1}], \quad (6)$$

where $^\vee$ denotes the p -adic completion. So $\mathcal{K}$ is a complete n -dimensional local field with residue field

$$k = O_{\mathcal{K}}/(p) \cong \mathbb{F}_p((T_1)) \dots ((T_{n-1})). \quad (7)$$

Remark. The topology on $\mathcal{K}$ is given by the p -adic valuation v .

We also define

$$A_1 = \mathbb{Z}_p[[T_1]][T_1^{-1}]^\vee, \quad (8)$$

$$A_i = A_{i-1}[[T_i]][T_i^{-1}]^\vee \text{ for } 2 \leq i < n, \quad (9)$$

$$O_{\mathcal{H}} = A_{n-1}, \quad (10)$$

$$\mathcal{H} = O_{\mathcal{H}}[p^{-1}], \quad (11)$$

so $\mathcal{K}$ is a finite extension of $\mathcal{H}$ of degree $[K : \mathbb{Q}_p]$. Note that the residue field h of $\mathcal{H}$ is isomorphic to k . Define the following towers of fields

$$H_k = \mathbb{Q}_p(\mu_{p^k}), \quad (12)$$

$$K_k = K(\mu_{p^k}), \quad (13)$$

$$\mathcal{H}_k = \mathcal{H}(\mu_{p^k}, T_1^{\frac{1}{p^k}}, \dots, T_n^{\frac{1}{p^k}}), \quad (14)$$

$$\mathcal{K}_k = \mathcal{K}(\mu_{p^k}, T_1^{\frac{1}{p^k}}, \dots, T_n^{\frac{1}{p^k}}), \quad (15)$$

and denote by $H_\infty, K_\infty, \mathcal{H}_\infty$ and $\mathcal{K}_\infty$ the respective direct limits. Note that $\mathcal{K}_\infty$ is a finite extension of $\mathcal{H}_\infty$ of degree $[K_\infty : H_\infty]$. For simplicity, we make the following assumption:-

Assumption. The fields H_∞ and K_∞ have the same residue field (which is isomorphic to $\mathbb{F}_p((T_1)) \dots ((T_{n-1}))$).

Let $\bar{\pi}_K = (\bar{\pi}_{K,n})_{n \geq 1}$ be a norm-compatible system of uniformizers of the tower $(K_n)_{n \geq 1}$.

Lemma 1. *The field of norms $\mathbb{E}_{\mathcal{K}}$ of $\mathcal{K}_\infty$ is given by*

$$\mathbb{E}_{\mathcal{K}} \cong (\mathbb{F}_p[[\bar{\pi}_K]] \hat{\otimes} \mathbb{F}_p((T_1)) \dots ((T_{n-1})))[\bar{\pi}_K^{-1}] \quad (16)$$

$$\cong \mathbb{F}_p((T_1)) \dots ((T_{n-1}))((\bar{\pi}_K)). \quad (17)$$

Proof. See the section on Kummer towers in [13]. (Note that $\mathbb{F}_p((T_1)) \dots ((T_{n-1}))$ has the discrete topology.) $\square$

Similarly, the field of norms of $\mathcal{H}$ is given by

$$\mathbb{E}_{\mathcal{H}} \cong \mathbb{F}_p((T_1)) \dots ((T_{n-1}))((\bar{\pi})), \quad (18)$$

where $\bar{\pi} = \epsilon - 1$ is the canonical system of uniformizers of the tower $(H_n)_{n \geq 1}$. (For the definition of ϵ , see e.g. [2].)

We also define

$$\mathbb{E}_{\mathcal{K}}^+ = \mathbb{F}_p((T_1)) \dots ((T_{n-1}))[[\bar{\pi}_K]], \quad (19)$$

$$\mathbb{E}_{\mathcal{H}}^+ = \mathbb{F}_p((T_1)) \dots ((T_{n-1}))[[\bar{\pi}]]. \quad (20)$$

2.2 Lift to characteristic 0

In this subsection, we define the higher dimensional analogues of Fontaine's rings $\mathbb{A}_{\mathbb{Q}_p}$ and $\mathbb{A}_K$. We start with the lift of $\mathbb{E}_{\mathcal{H}}$:-

Definition. Let $\pi = [\epsilon] - 1$ be the usual lift of $\bar{\pi}$ to characteristic 0 (where $[\epsilon]$ denotes the Teichmueller representative of ϵ). Let $\mathcal{R}_{\mathbb{Q}_p}^+ = \mathcal{O}_{\mathcal{H}}[[\pi]]$ and $\mathcal{R}_{\mathbb{Q}_p} = \mathcal{R}_{\mathbb{Q}_p}^+[\pi^{-1}]^\vee$, so $\mathcal{R}_{\mathbb{Q}_p}$ is a complete regular local ring of dimension n with residue field $\mathcal{R}_{\mathbb{Q}_p}/(p) \cong \mathbb{E}_{\mathcal{H}}$.

Note. $\mathbb{A}_{\mathbb{Q}_p} \subset \mathcal{R}_{\mathbb{Q}_p}$ and $\mathbb{A}_{\mathbb{Q}_p}^+ \subset \mathcal{R}_{\mathbb{Q}_p}^+$.

Action of Frobenius. The action of the Frobenius operator ϕ on $\mathcal{R}_{\mathbb{Q}_p}$ is determined by

$$\phi : T_i \rightarrow T_i^p \text{ for all } i \quad (21)$$

$$\pi \rightarrow (\pi + 1)^p - 1. \quad (22)$$

Lemma 2. *The ring $\mathcal{R}_{\mathbb{Q}_p}$ is a free $\phi(\mathcal{R}_{\mathbb{Q}_p})$ -module of rank p^n , and a basis is given by $\{T_1^{i_1} \dots T_{n-1}^{i_{n-1}} (\pi + 1)^j\}_{0 \leq i_1, \dots, i_{n-1}, j \leq p-1}$.*

Proof. We can assume that the action of ϕ on T_i is trivial for all i . The lemma then follows by explicit calculation. $\square$

Construction of the lift $\mathcal{R}_K$ of $\mathbb{E}_K$. We start by proving the following lemma:-

Lemma 3. *The field $\mathbb{E}_K$ is given by*

$$\mathbb{E}_K \cong \mathbb{E}_K \otimes_{\mathbb{E}_H} \mathbb{E}_{\mathcal{H}} \quad (23)$$

$$= \mathbb{F}_p((\bar{\pi}_K)) \otimes_{\mathbb{F}_p((\bar{\pi}))} \mathbb{F}_p((X_1)) \dots ((X_{n-1}))((\bar{\pi})) \quad (24)$$

Proof. We have $[\mathbb{E}_K : \mathbb{E}_{\mathcal{H}}] = [\mathbb{E}_K : \mathbb{E}_H] = [K_\infty : H_\infty]$, and there is a natural injection

$$\mathbb{F}_p((\bar{\pi}_K)) \otimes_{\mathbb{F}_p((\bar{\pi}))} \mathbb{F}_p((X_1)) \dots ((X_{n-1}))((\bar{\pi})) \hookrightarrow \mathbb{F}_p((X_1)) \dots ((X_{n-1}))((\bar{\pi}_K)), \quad (25)$$

given by 'component-wise' multiplication. But $\mathbb{F}_p((\bar{\pi}_K))$ is an $\mathbb{F}_p((\bar{\pi}))$ -vector space of dimension $[K_\infty : H_\infty]$, so

$$\dim_{\mathbb{E}_{\mathcal{H}}} \mathbb{F}_p((\bar{\pi})) \otimes_{\mathbb{F}_p((\bar{\pi}))} \mathbb{E}_{\mathcal{H}} = [K_\infty : H_\infty]. \quad (26)$$

$\square$

We can now make the following definitions:-

Definition. Let $\mathcal{R}_K = \mathbb{A}_K \otimes_{\mathbb{A}_{\mathbb{Q}_p}} \mathcal{R}_{\mathbb{Q}_p}$, and let $\mathcal{S}_K = \mathcal{R}_K[p^{-1}]$.

Then $\mathcal{S}_K$ is an $n + 1$ -dimensional local field (in the p -adic topology) and $\mathcal{R}_K$ is a p -adically complete local ring of dimension $n + 1$ and residue field $\mathbb{E}_K$, so it is a lift of $\mathbb{E}_K$ to characteristic 0. The natural topology on $\mathcal{R}_K$ is

weaker than the p -adic topology:- A family of neighbourhoods of 0 in $\mathcal{R}_K$ is given by $p^k \mathcal{R}_K + \pi_K^l O_{\mathcal{H}}[[\pi_K]]$ for $k, l \geq 1$. (One can show that this definition is independent of the lift π_K of $\bar{\pi}_K$.) The natural topology on $\mathcal{S}_K$ is then the inductive limit topology.

The action of the Frobenius operator ϕ extends naturally to $\mathcal{R}_K$ and $\mathcal{S}_K$, with the usual action on $\mathbb{A}_K$. Note that the action of ϕ restricts to $\mathcal{R}_K^+ = \{a \in \mathcal{R}_K : a \bmod p \in \mathbb{E}_K^+\}$.

Lemma 4. $\mathcal{R}_K^+$ is a free $\phi(\mathcal{R}_K^+)$ -module of rank p^n .

Proof. It follows from the classical theory that $\mathbb{A}_K^+$ is a free $\phi(\mathbb{A}_K^+)$ -module of dimension p with basis $1, \pi + 1, \dots, (\pi + 1)^{p-1}$, and $\mathcal{R}_{\mathbb{Q}_p}^+$ is a free $\phi(\mathcal{R}_{\mathbb{Q}_p}^+)$ -module of dimension p^n with basis $\{T_1^{i_1} \dots T_{n-1}^{i_{n-1}} (\pi + 1)^j\}_{0 \leq i_1, \dots, i_{n-1}, j \leq p-1}$. $\square$

Similarly, we have

Lemma 5. $\mathcal{R}_K$ is a free $\phi(\mathcal{R}_K)$ -module of rank p^n , and $\mathcal{S}_K$ is a $\phi(\mathcal{S}_K)$ -vector space of dimension p^n .

2.3 Some details about $\phi(\pi_K)$

Recall the following result from [1]:-

Proposition 6. Let $e = [K : \mathbb{Q}_p]$. Then there exists $n(K) \in \mathbb{N}$ and an element $\pi_K \in \mathbb{A}_K^{\dagger, n(K)}$ such that the image of $\pi_K \bmod p$ is the uniformizing element $\bar{\pi}_K$ of $\mathbb{E}_K$. Furthermore, if $n \geq n(K)$, then $\mathbb{A}_K^{\dagger, n}$ is isomorphic to the ring of series $\sum_{k \in \mathbb{Z}} c_k X^k$, $c_k \in \mathbb{Z}_p$ for all k , which are analytic and bounded above by 1 on the annulus $\{p^{-\frac{1}{e(p-1)p^{n-1}}} \leq |X| < 1\}$.

Remark. This result is hidden in the proof of Proposition III.2.1 in [3] - the main point is that for $n \gg 0$, we have $\mathbb{A}_K^{\dagger, n} = \mathbb{A}_{\mathbb{Q}_p}^{\dagger, n}[\pi_K]$, where $\pi_K \in \mathbb{A}_K$ is a lift of $\bar{\pi}_K$.

Proposition 6 has the following useful consequence:-

Corollary 7. We have that $\mathbb{A}_K = \mathbb{A}_{\mathbb{Q}_p}[\pi_K]$. It follows that any element of $\mathbb{A}_K$ - and hence any element of $\mathcal{R}_K$ - can be written as a power series in π_K .

It is shown in [2] that the power series from Proposition 6 can be characterized as follows:-

Lemma 8. Let $x = \sum_{k \in \mathbb{Z}} c_k X^k$ with $c_k \in \mathbb{Z}_p$ for all k . Then x is analytic and bounded above by 1 on the annulus $\{p^{-\frac{1}{e(p-1)p^{n-1}}} \leq |X| < 1\}$ if and only if $v_p(c_k) + \frac{k}{(p-1)p^{n-1}} \geq 0$ for all $k < 0$ and $\rightarrow +\infty$ as $k \rightarrow -\infty$.

Corollary 9. Let $n \geq n(K)$, and let $x = \sum_{k \in \mathbb{Z}} c_k \pi_K^k \in \mathbb{A}_K^{\dagger, n}$. If $c_0 \in \mathbb{Z}_p^\times$, then x is invertible in $\mathbb{A}_K^{\dagger, n+1}$.

For the rest of this section we fix this lift π_K and assume that $n > n(K)$. Our aim is to prove Corollary 13 below, which gives some details about the action of ϕ on π_K . Recall that by our simplifying assumption $e = [K_\infty : F_\infty]$.

To prove Corollary 13, we first need a couple of lemmas:-

Lemma 10. *Let $n \gg 0$, and let $x \in \mathbb{A}_K^{\dagger, n}$. Write x as a power series in π_K , i.e.*

$$x = \cdots + c_{-2}\pi_K^{-2} + c_{-1}\pi_K^{-1} + c_0 + c_1\pi_K + \cdots, \quad (27)$$

and let

$$x' = \cdots - 2c_{-2}\pi_K^{-3} - c_{-1}\pi_K^{-2} + c_1 + 2c_2\pi_K + \cdots \quad (28)$$

be the formal derivative $x' = \frac{dx}{d\pi_K}$. Then $x' \in \mathbb{A}_K^{\dagger, n}$.

Proof. By the definition of $\mathbb{A}_K^{\dagger, n}$, we know that

$$v_p(c_i) + \frac{i}{(p-1)ep^{n-1}} \geq 0 \text{ for all } i < 0 \quad (29)$$

$$v_p(c_i) + \frac{i}{(p-1)ep^{n-1}} \rightarrow +\infty \text{ as } i \rightarrow -\infty \quad (30)$$

Now $v_p(c_i)$ is always a positive integer, so if $((p-1)ep^{n-1}) \nmid i$, then

$$v_p(c_i) + \frac{i}{(p-1)ep^{n-1}} \geq \frac{1}{(p-1)ep^{n-1}} > 0, \quad (31)$$

so certainly

$$v_p(ic_i) + \frac{i-1}{(p-1)ep^{n-1}} \geq 0. \quad (32)$$

Now if $((p-1)ep^{n-1}) \mid i$, then $v_p(i) \geq n-1$, so

$$v_p(ic_i) + \frac{i-1}{(p-1)ep^{n-1}} \geq 0. \quad (33)$$

Also, since

$$v_p(c_i) + \frac{i}{(p-1)ep^{n-1}} \rightarrow +\infty \text{ as } i \rightarrow -\infty, \quad (34)$$

we see that

$$v_p(c_i) + \frac{i-1}{(p-1)ep^{n-1}} \rightarrow +\infty \text{ as } i \rightarrow -\infty, \quad (35)$$

so certainly

$$v_p(ic_i) + \frac{i-1}{(p-1)ep^{n-1}} \rightarrow +\infty \text{ as } i \rightarrow -\infty, \quad (36)$$

which finishes the proof. $\square$

Lemma 11. *Let $x \in \mathbb{A}_K^{\dagger, N}$ for some $N \gg 0$. Write*

$$x = \cdots + a_{-1}\pi_K^{-1} + a_0 + a_1\pi_K + a_2\pi_K^2 + \cdots,$$

and assume that x is precisely divisible by p^k in $\mathbb{A}_K$. Then there exists $M > 0$ and $L \geq 0$ such that

$$x = p^k \pi_K^{-L} y$$

for some $y \in (\mathbb{A}_K^{\dagger, N+M})^\times$.

Proof. Let L be minimal such that $v_p(a_L) = k$. Suppose first that $L = 0$. Now

$$v_p(a_i) - k + \frac{i}{(p-1)ep^{N-1}} \rightarrow +\infty \text{ when } i \rightarrow -\infty,$$

so there exists $K \gg 0$ such that $v_p(a_i) - k + \frac{i}{(p-1)ep^{N-1}} \geq 0$ for all $i \leq -K$. Choose $M > 0$ such that

$$\frac{K}{(p-1)ep^{N+M-1}} < 1.$$

We can procede analogously if $L < 0$. If $L > 0$, then choose M such that $p^M > L + 1$. Since by the choice of L the constant term of the π_K -expansion will be in $\mathbb{Z}_p^\times$, it follows from Corollary 9 that y will be invertible in $\mathbb{A}_K^{\dagger, N+M}$ if M is sufficiently large. $\square$

We assume for now that $p \mid e$. (We can argue similarly when e and p are coprime.) By the theory of (1-dimensional) fields of norms, $\mathbb{E}_K$ is a separable extension of $\mathbb{E}_{\mathbb{Q}_p}$ of degree e , so since $\bar{\pi}_K$ is a uniformizer of $\mathbb{E}_K$, it satisfies a separable polynomial

$$\bar{f}(X) = X^e + \bar{a}_{e-1}X^{e-1} + \cdots + \bar{a}_0, \quad (37)$$

where $\bar{a}_i \in \mathbb{E}_{\mathbb{Q}_p}^+$ for all i . Since $\bar{f}(X)$ is separable, there exists $0 \leq i \leq e$ such that $p \nmid i$ and $\bar{a}_i \neq 0$. Now $\bar{a}_0$ and $\bar{\pi}$ have the same valuation in $\mathbb{E}_K$, so $\bar{a}_0 = \bar{\pi} \times a$, where $a \in 1 + \bar{\pi}_K \mathbb{E}_K^+$ (or maybe multiplied by an element in $\mathbb{F}_p^\times$, but we ignore that). Using (37), one can now expand $\bar{\pi}$ as a power series in $\bar{\pi}_K$,

$$\bar{\pi} = \bar{\pi}_K^e (1 + \bar{b}_1 \bar{\pi}_K + \bar{b}_2 \bar{\pi}_K^2 + \cdots), \quad (38)$$

where $\bar{b}_i \in \mathbb{F}_p$. By the remark following (37), there exists $i > 0$ such that $\bar{b}_i \neq 0$ and $p \nmid i$. Let I be the minimal such integer.

Now $\pi_K \in \mathbb{A}_K$ is a lift of $\bar{\pi}_K$, so it satisfies a polynomial

$$f(X) = X^e + a_{e-1}X^{e-1} + \cdots + a_0, \quad (39)$$

where $a_i \in \mathbb{A}_{\mathbb{Q}_p}$ and $f(X) \bmod p = \bar{f}(X)$. In fact, it is shown in the proof of Proposition III.2.1 in [3] that $a_i \in \mathbb{A}_{\mathbb{Q}_p}^+$. As above, we can use equation (39) to expand π as a power series in π_K ,

$$\pi = \pi_K^e (b_0 + b_1 \pi_K + b_2 \pi_K^2 + \cdots), \quad (40)$$

which reduces to (38) when taken $\pmod p$, i.e.

$$b_0 = 1 \pmod p, \quad (41)$$

$$b_i \pmod p = \bar{b}_i \quad \text{for all } i \geq 1. \quad (42)$$

Applying ϕ to (40) gives

$$(\pi + 1)^p - 1 = \phi(\pi_K)^e (b_0 + b_1 \phi(\pi_K) + b_2 \phi(\pi_K)^2 + \dots). \quad (43)$$

Write

$$\phi(\pi_K) = \dots + c_{-1} \pi_K^{-1} + c_0 + c_1 \pi_K + c_2 \pi_K^2 + \dots$$

with $c_i \in \mathbb{Z}_p$. (Such an expansion exists since $\phi(\pi_K) \in \mathbb{A}_K^{\dagger, n}$ for all $n \geq 2$, and by 6 all elements of $\mathbb{A}_K^{\dagger, n}$ can be written as power series in π_K for all $n \gg 0$.)

Proposition 12. *There exists $i \leq -1$ such that $p \nmid i$, $p \mid c_i$ but $p^2 \nmid c_i$.*

Proof. Recall that I is the minimal positive integer not divisible by p such that $p \nmid b_I$. Expanding the left hand side of (43) as a power series in π_K shows that the coefficient of π_K^I is precisely divisible by p . Note that $\phi(\pi_K) \pmod p = \bar{\pi}_K^p$, i.e. $p \mid c_i$ for all $i \neq p$. Since $p \mid e$, it follows that in the power series expansion of $\phi(\pi_K)^e$, every coefficient apart from the one of π_K^{pe} is divisible by p^2 . Now if $p \mid b_j$, then it is clear that the coefficient of π_K^I in $b_j \phi(\pi_K)^{e+j}$ is divisible by p^2 . (Recall that $p \mid c_i$ for all $i \neq p$.) Similarly, if j is a positive integer divisible by p , then the coefficient of π_K^I in $b_j \phi(\pi_K)^{e+j}$ is divisible by p^2 . Recall that by the choice of I , $p \mid b_j$ for all $0 < j < I$. Since in the power series expansion of $\phi(\pi_K)^e$ every coefficient apart from the one of π_K^{pe} is divisible by p^2 , there must exist some $j \geq I$ not divisible by p such that $c_{-p(e+j-1)+e+I}$ is precisely divisible by p . Now by assumption $e \geq p$ and $p \neq 2$, which implies that $p(e+j-1) > e+I$ and finishes the proof. $\square$

Corollary 13. *Let $\phi(\pi_K) = \dots + c_{-1} \pi_K^{-1} + c_0 + c_1 \pi_K + c_2 \pi_K^2 + \dots$, where $c_i \in \mathbb{Z}_p$ for all i , and let $g(\pi_K) = \frac{d}{d\pi_K} \phi(\pi_K)$ be its formal derivative. Then there exists $I < 0$ such that*

$$g(\pi_K) = p \pi_K^I \times a, \quad (44)$$

where $p \nmid I$ and $a \in (\mathbb{A}_K^{\dagger, n})^\times$ for some $n \gg 0$.

Proof. Let $I < 0$ be minimal such that $p \nmid I$ and c_I is precisely divisible by p (i.e. not divisible by p^2). Then we can write

$$g(\pi_K) = p \pi_K^I a, \quad (45)$$

and since $p \nmid I$, we have $a \in (\mathbb{A}_K^{\dagger, n})^\times$ for $n \gg 0$ by Lemma 11. $\square$

3 Some stuff we need

3.1 Milnor K -groups and completions

Notation. For any commutative ring A , denote by $K_n(A)$ Milnor's K_n -group of A .

Recall that $K_n(A)$ is the group generated by symbols $\{a_1, \dots, a_n\}$, $a_i \in A^\times$ for all i , subject to the relations

- (1) $\{a_1, \dots, a_n\}$ is multiplicative in each a_i ;
- (2) $\{a_1, \dots, a_n\} = 1$ if $a_i + a_j = 1$ for some $i \neq j$.

It follows from these relations that interchanging any two entries of $a = \{a_1, \dots, a_n\}$ yields the inverse a^{-1} . In particular, if $a_i = a_j$ for some $i \neq j$, then $a^2 = 1$.

Definition. Let U be the subgroup of $\mathcal{R}_K$ consisting of the elements x such that $x \bmod p \in \mathbb{E}_K^+$, and let $\mathcal{U}$ be the subgroup of $K_n(\mathcal{S}_K)$ generated by the symbols $\{a_1, \dots, a_n\}$ with $a_1 \in U^\times$, $a_i \in \mathcal{S}_K^\times$. Define

$$\tilde{K}_n(\mathcal{S}_K) = \varprojlim K_n(\mathcal{S}_K)/p^m \mathcal{U}. \quad (46)$$

Definition. Let $\mathcal{V}$ be the subgroup of $K_n(\mathbb{E}_K)$ generated by the symbols $\{a_1, \dots, a_n\}$ with $a_1 \in 1 + \mathbb{E}_K^+$, $a_i \in \mathbb{E}_K^\times$. Define

$$\tilde{K}_n(\mathbb{E}_K) = \varprojlim K_n(\mathbb{E}_K)/p^m \mathcal{V}. \quad (47)$$

When A is a local field with maximal ideal $\mathfrak{m}$, we also define the $\mathfrak{m}$ -adic completion $\hat{K}_n(A)$ of $K_n(A)$ as follows:-

Definition. For all $r \geq 1$ let $\mathcal{U}^{(r)}$ be the subgroup of $K_n(A)$ generated by the symbols of the form $\{a_1, \dots, a_n\}$, where $a_1 \in 1 + \mathfrak{m}^r$ and $a_i \in A^\times$ for all i . Then define

$$\hat{K}_n(A) = \varprojlim K_n(A)/\mathcal{U}^{(r)}. \quad (48)$$

In particular, we will be interested in the case when $A = \mathbb{E}_K$:-

Lemma 14. *We have a continuous map*

$$\tilde{K}_n(\mathbb{E}_K) \longrightarrow \hat{K}_n(\mathbb{E}_K).$$

Proof. Let $U^{(r)} = 1 + \pi_K^r \mathbb{E}_K^+$, and let $\mathcal{U}^{(r)}$ be the subgroup of $K_n(\mathbb{E}_K)$ generated by symbols of the form $\{a_1, \dots, a_n\}$ where $a_1 \in U^{(r)}$ and $a_i \neq 0$. It follows from the definitions of the filtrations that for all $k \geq 1$ we have $p^k K_n(\mathbb{E}_K) \subset \mathcal{U}^{(k)}$. $\square$

Note. The above map has dense image.

3.2 Modules of differentials

Definition. Let $\Omega_{\mathcal{R}_K/\mathbb{Z}}^{n-1}$ be the module of absolute differential $(n-1)$ -forms of $\mathcal{R}_K$, and define $\Omega_{\mathcal{R}_K}^{n-1}$ to be its p -adic completion

$$\Omega_{\mathcal{R}_K}^{n-1} = \varprojlim \Omega_{\mathcal{R}_K/\mathbb{Z}}^{n-1}/p^m \Omega_{\mathcal{R}_K/\mathbb{Z}}^{n-1}. \quad (49)$$

Lemma 15. *We have an isomorphism of $\mathcal{R}_K$ -modules*

$$\Omega_{\mathcal{R}_K}^{n-1} \cong \bigoplus \mathcal{R}_K d\log(\pi_K) \wedge d\log(T_{i_1}) \wedge \cdots \wedge d\log(T_{i_{n-2}}) \oplus \mathcal{R}_K \bigwedge_{1 \leq i < n} d\log(T_i) \quad (50)$$

Proof. Follows from explicit computation, using the observation that $\pi_K + 1$ is invertible in $\mathcal{R}_K^+$. (Note that $d\log(\pi_K) = \frac{\pi_K+1}{\pi_K} d\log(\pi_K + 1)$.) $\square$

Now $\mathcal{R}_K$ is finite flat and locally of complete intersection over $\phi(\mathcal{R}_K)$, so as shown in [7], we have a trace map of differential forms

$$\mathrm{Tr}_\phi^{(n-1)} : \Omega_{\mathcal{R}_K}^{n-1} \longrightarrow \Omega_{\mathcal{R}_K}^{n-1} \quad (51)$$

and of course the usual trace map

$$\mathrm{Tr}_\phi : \mathcal{R}_K \longrightarrow \mathcal{R}_K. \quad (52)$$

Lemma 16. *The trace map is characterized as follows:-*

$$\begin{aligned} \mathrm{Tr}_\phi^{(n-1)}(\alpha \cdot \bigwedge d\log(T_{i_j}) \wedge d\log(\pi_K + 1)) &= \frac{1}{p^{n-1}} \mathrm{Tr}_\phi(p \cdot \alpha \omega) \bigwedge d\log(T_{i_j}) \wedge d\log(\pi_K + 1), \\ \mathrm{Tr}_\phi^{(n-1)}(\beta d\log(T_1) \wedge \cdots \wedge d\log(T_{n-1})) &= \frac{1}{p^{n-1}} \mathrm{Tr}_\phi(\beta) d\log(T_1) \wedge \cdots \wedge d\log(T_{n-1}), \end{aligned}$$

where

$$\omega = \frac{\phi(\pi_K) + 1}{\pi_K + 1} \left(\frac{d}{d\pi_K} \phi(\pi_K + 1) \right)^{-1}. \quad (53)$$

Proof. First note that since ϕ is a ring homomorphism, we have $\frac{d}{d\pi_K} \phi(\pi_K + 1) = \frac{d}{d\pi_K} \phi(\pi_K)$. Now by Corollary 13, there exists $I < 0$ and $a \in (\mathbb{A}_K^{\dagger, N})^\times$ for $N \gg 0$ such that

$$\frac{d}{d\pi_K} \phi(\pi_K) = p\pi_K^I a,$$

so $p\omega \in \mathcal{S}_K^{\dagger, N}$ for $N \gg 0$. The lemma is now an immediate consequence of Remark (iii) in Section 2.1 in [7]. $\square$

The following corollary will be important later:-

Corollary 17. $\mathrm{Tr}_\phi^{(n-1)}(\Omega_{\mathcal{R}_K}^{n-1}) \subset p\Omega_{\mathcal{R}_K}^{n-1}$

Proof. This follows from the equations in Lemma 16 and the observation that $\mathrm{Tr}_\phi(\Omega_{\mathcal{R}_K}^{n-1}) \subset p^n \Omega_{\mathcal{R}_K}^{n-1}$. $\square$

3.3 The norm map

It follows from Milnor K -theory that the p -adic valuation on $\mathcal{S}_K$ induces a surjective map

$$\lambda : K_n(\mathcal{S}_K) \longrightarrow K_n(\mathbb{E}_K) \quad (54)$$

satisfying the following condition:- If $a_i \in \mathcal{R}_K^\times$ and $\bar{a}_i$ denotes the image of a_i in $\mathbb{E}_K$, then

$$\lambda : \{a_1 p^{i_1}, \dots, a_n p^{i_n}\} \longrightarrow \{\bar{a}_1, \dots, \bar{a}_n\}. \quad (55)$$

Clearly λ extends to a map on the p -adic completions

$$\lambda : \tilde{K}_n(\mathcal{S}_K) \longrightarrow \tilde{K}_n(\mathbb{E}_K). \quad (56)$$

Note. Every element in $\tilde{K}_n(\mathcal{S}_K)$ can be written as the (infinite) product of symbols $\{a_1, \dots, a_n\}$ where each a_i either lies in $\mathcal{R}_K^\times$ or is equal to p .

Let $\mathcal{V}$ be the subgroup of $\tilde{K}_n(\mathcal{S}_K)$ topologically generated by the symbols $\{a_1, \dots, a_n\}$, where at least one a_i is equal to p . Note that by the definition of λ , we have $\mathcal{V} \subset \ker(\lambda)$. Equip $\tilde{K}_n(\mathcal{S}_K)/\mathcal{V}$ with the product topology.

Now $\mathcal{S}_K$ is a finite dimensional vector space over $\phi(\mathcal{S}_K)$ of dimension p^n (and $\mathbb{E}_K$ is a finite dimensional vector space of $\phi(\mathbb{E}_K)$ of dimension p^n), so as shown in Corollary 7.6.3 in [14] we have the following result:-

Lemma 18. *We have a commutative diagram*

$$\begin{array}{ccc} K_n(\mathcal{S}_K) & \xrightarrow{\lambda} & K_n(\mathbb{E}_K) \\ N \downarrow & & \downarrow N \\ K_n(\mathcal{S}_K) & \xrightarrow{\lambda} & K_n(\mathbb{E}_K) \end{array}$$

where N are the respective norm maps induced from the finite field extensions $\mathcal{S}_K/\phi(\mathcal{S}_K)$ and $\mathbb{E}_K/\phi(\mathbb{E}_K)$.

Note. Since the norm maps are group homomorphisms, the above diagram extends to the completed K -groups $\tilde{K}_n(\mathcal{S}_K)$ and $\tilde{K}_n(\mathbb{E}_K)$.

Lemma 19. *The action of N factors through $\mathcal{V}$.*

Proof. Clear from the projection formula. □

We want to prove the following result:-

Proposition 20. *The map λ gives a canonical isomorphism*

$$(\tilde{K}_n(\mathcal{S}_K)/\mathcal{V})^{N=1} \cong \tilde{K}_n(\mathbb{E}_K). \quad (57)$$

Definition. Let $\mathcal{U}$ be the subgroup of $\tilde{K}(\mathcal{S}_K)/\mathcal{V}$ topologically generated by the symbols of the form $\{a_1, \dots, a_n\}$, where $a_1 \in 1 + p\mathcal{R}_K$ and $a_i \in \mathcal{R}_K^\times$ for all i .

Lemma 21. *We have a short exact sequence*

$$1 \longrightarrow \mathcal{U} \longrightarrow \tilde{K}_n(\mathcal{S}_K)/\mathcal{V} \longrightarrow \tilde{K}_n(\mathbb{E}_K) \longrightarrow 1. \quad (58)$$

Proof. Let $x \in \tilde{K}_n(\mathcal{S}_K)$. Then there exist $x_i \in K_n(\mathcal{S}_K)$ for $i \geq 0$ such that x can be represented by the infinite product $\prod_i x_i^{p^i}$. Now each of the x_i can be written as a product of symbols $\{a_1^{(i)}, \dots, a_n^{(i)}\}$ where either one of the $a_j^{(i)}$ is equal to p or all of the $a_j^{(i)}$ are invertible in $\mathcal{R}_K$. It follows that the natural map $\tilde{K}_n(\mathcal{R}_K) \rightarrow \tilde{K}_n(\mathcal{S}_K)$ induces a surjection $\tilde{K}_n(\mathcal{R}_K) \rightarrow \tilde{K}_n(\mathcal{S}_K)/\mathcal{V}$, and we have a commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathcal{U} & \longrightarrow & \tilde{K}_n(\mathcal{S}_K)/\mathcal{V} & \longrightarrow & \tilde{K}_n(\mathbb{E}_K) \longrightarrow 1 \\ & & \uparrow \alpha & & \uparrow \beta & & \uparrow \gamma \\ 1 & \longrightarrow & \mathcal{W} & \longrightarrow & \tilde{K}_n(\mathcal{R}_K) & \longrightarrow & \tilde{K}_n(\mathbb{E}_K) \longrightarrow 1 \end{array}$$

Now γ is the identity, so α is surjective by the snake lemma. It is therefore sufficient to show that $\mathcal{W}$ is topologically generated by symbols of the form $\{a_1, \dots, a_n\}$, where $a_1 \in 1 + p\mathcal{R}_K$ and $a_i \in \mathcal{R}_K^\times$ for all i . Now we have a short exact sequence

$$1 \longrightarrow W \longrightarrow K_n(\mathcal{R}_K) \longrightarrow K_n(\mathbb{E}_K) \longrightarrow 1, \quad (59)$$

and imitating the proof of Proposition 4.1 in [8] shows that W is the subgroup generated by symbols of the form $\{a_1, \dots, a_n\}$, where $a_1 \in 1 + p\mathcal{R}_K$ and $a_i \in \mathcal{R}_K^\times$ for all i . Passing to the completions of (59) gives the bottom row of the above diagram and hence finishes the proof. $\square$

Lemma 22. *The action of N on $\tilde{K}_n(\mathbb{E}_K)$ is trivial.*

Proof. As $\phi : \mathbb{E}_K \rightarrow \mathbb{E}_K$ is the p th power map, the result follows by applying Lemma 12, paragraph 3.3 in [9]. $\square$

In view of the preceding two lemmas, to prove Proposition 20 it is sufficient to prove the following result:-

Lemma 23. *The action of N on $\mathcal{U}$ is nilpotent.*

Proof. Let

$$\exp_{p,n-1} : \Omega_{\mathcal{R}_K}^{n-1} \longrightarrow \mathcal{U} \quad (60)$$

be the exponential homomorphism for the K_n -group defined by Kurihara in [12] which is characterised as follows:- For $\alpha \in \mathcal{R}_K$, $\beta_1, \dots, \beta_{n-1} \in \mathcal{R}_K^\times$,

$$\exp_{p,n-1}(\alpha \cdot d \log(\beta_1) \wedge \dots \wedge d \log(\beta_{n-1})) = \{\exp(p\alpha), \beta_1, \dots, \beta_{n-1}\} \quad (61)$$

Since $\Omega_{\mathcal{R}_K}^{n-1}$ is p -adically complete, one can check that the map is surjective. (Note that $\exp_{p,n-1}$ is continuous.) The equations in Lemma 16 show that we have a commutative diagram

$$\begin{array}{ccc} \exp_{p,n-1} : \Omega_{\mathcal{R}_K}^{n-1} & \longrightarrow & \tilde{K}_n(\mathcal{S}_K)/\mathcal{V} \\ \downarrow \text{Tr}_\phi^{(n-1)} & & \downarrow N \\ \exp_{p,n-1} : \Omega_{\mathcal{R}_K}^{n-1} & \longrightarrow & \tilde{K}_n(\mathcal{S}_K)/\mathcal{V} \end{array}$$

Now Corollary 17 shows that $\text{Tr}_\phi^{(n-1)}$ is nilpotent on $\Omega_{\mathcal{R}_K}^{n-1}$, which finishes the proof. $\square$

4 The logarithmic derivative

In this section, we will use Proposition 20 to construct the (generalized) logarithmic derivative. But before we do this, we need to introduce another module of differentials:-

4.1 Some more modules of differentials

Definition. Let $\Omega_{\mathcal{R}_K/\mathbb{Z}}^1$ be the module of absolute differential 1-forms of $\mathcal{R}_K$, and let

$$\Omega_{\mathcal{R}_K^+/\mathbb{Z}}^1(\log) = (\Omega_{\mathcal{R}_K^+/\mathbb{Z}}^1 \oplus \mathcal{R}_K^+ \otimes_{\mathbb{Z}} \mathcal{R}_K^\times) / \mathcal{N}, \quad (62)$$

where $\mathcal{N}$ is the $\mathcal{R}_K^+$ -submodule of the direct sum generated by the elements $(-da, a \otimes a)$ for $a \in \mathcal{R}_K^+ \cap \mathcal{R}_K^\times$. In $\Omega_{\mathcal{R}_K^+/\mathbb{Z}}^1(\log)$, denote the class $(0, 1 \otimes a)$ for $a \in \mathcal{S}_K^\times$ by $d \log(a)$. Let $\Omega_{\mathcal{R}_K^+}^n(\log)$ be its p -adic completion

$$\Omega_{\mathcal{R}_K^+}^n(\log) = \varprojlim \Omega_{\mathcal{R}_K^+/\mathbb{Z}}^n(\log) / p^m \Omega_{\mathcal{R}_K^+/\mathbb{Z}}^n(\log). \quad (63)$$

Similarly, define

$$\Omega_{\mathcal{R}_K^+/\mathbb{Z}}^1(\log)(\log) = (\Omega_{\mathcal{R}_K^+/\mathbb{Z}}^1 \oplus \mathcal{R}_K^+ \otimes_{\mathbb{Z}} \mathcal{S}_K^\times) / \mathcal{M}, \quad (64)$$

where $\mathcal{M}$ is the $\mathcal{R}_K^+$ -submodule of the direct sum generated by the elements $(-da, a \otimes a)$ for $a \in \mathcal{R}_K^+ \cap \mathcal{S}_K^\times$. In $\Omega_{\mathcal{R}_K^+/\mathbb{Z}}^1(\log)(\log)$, denote the class $(0, 1 \otimes a)$ for $a \in \mathcal{S}_K^\times$ by $d \log(a)$. Let $\Omega_{\mathcal{R}_K^+}^n(\log)(\log)$ be its p -adic completion

$$\Omega_{\mathcal{R}_K^+}^n(\log)(\log) = \varprojlim \Omega_{\mathcal{R}_K^+/\mathbb{Z}}^n(\log)(\log) / p^m \Omega_{\mathcal{R}_K^+/\mathbb{Z}}^n(\log)(\log). \quad (65)$$

Finally, let

$$\Omega_{\mathcal{R}_K^+}^n = \varprojlim \Omega_{\mathcal{R}_K^+/\mathbb{Z}}^n / p^m \Omega_{\mathcal{R}_K^+/\mathbb{Z}}^n. \quad (66)$$

Lemma 24. *We have isomorphisms of $\mathcal{R}_K^+$ -modules*

$$\Omega_{\mathcal{R}_K^+}^n \cong \mathcal{R}_K^+ d\log(\pi_K + 1) \wedge d\log(T_{i_1}) \wedge \cdots \wedge d\log(T_{i_{n-1}}), \quad (67)$$

$$\Omega_{\mathcal{R}_K^+}^n(\log) \cong \mathcal{R}_K^+ d\log(\pi_K) \wedge d\log(T_{i_1}) \wedge \cdots \wedge d\log(T_{i_{n-1}}). \quad (68)$$

Proof. Follows from explicit computation, using the observation that $\pi_K + 1$ is invertible in $\mathcal{R}_K^+$. (Note that $d\log(\pi_K) = \frac{\pi_K + 1}{\pi_K} d\log(\pi_K + 1)$.) $\square$

Lemma 25. *The natural map $\Omega_{\mathcal{R}_K^+/\mathbb{Z}}^n(\log) \rightarrow \Omega_{\mathcal{R}_K^+/\mathbb{Z}}^n(\log)(\log)$ is injective, and its cokernel is killed by p .*

Proof. Let $a \in \mathcal{R}_K^+ \cap \mathcal{S}_K^\times$, and write $a = p^k b$, where $v(a) = 0$. Then

$$(-da, a \otimes a) = p^n(-db, b \otimes b) + b(0, p^n \otimes p^n). \quad (69)$$

Now $(0, p^n \otimes p^n) = -dp^n = 0$ and $p^n(-db, b \otimes b) \in \mathcal{N}$, so $(-da, a \otimes a)$ is already trivial in $\Omega_{\mathcal{R}_K^+}^n$. It follows that the above map is injective. Since $p(x, y \otimes p) = 0$ for all x, y , it follows that the cokernel of the map is killed by p . $\square$

Corollary 26. *Let U be the subgroup of $\Omega_{\mathcal{R}_K^+}^n(\log)(\log)$ generated by the elements of the form $\beta.d\log(p) \wedge d\log(b_2) \wedge \cdots \wedge d\log(b_n)$. We have a canonical isomorphism of $\mathcal{R}_K^+$ -modules*

$$\Omega_{\mathcal{R}_K^+}^n(\log)(\log)/U \cong \Omega_{\mathcal{R}_K^+}^n(\log). \quad (70)$$

We use this isomorphism to identify $\Omega_{\mathcal{R}_K^+}^n(\log)(\log)/U$ with $\Omega_{\mathcal{R}_K^+}^n(\log)$.

4.2 The ψ -operator

The module $\mathcal{R}_K^+$ is finite flat and locally of complete intersection over $\phi(\mathcal{R}_K^+)$, so as shown in [7], we have a trace map of differential forms

$$\mathrm{Tr}_\phi^{(n)} : \Omega_{\mathcal{R}_K^+}^n \longrightarrow \Omega_{\mathcal{R}_K^+}^n \quad (71)$$

and of course the usual trace map

$$\mathrm{Tr}_\phi : \mathcal{R}_K^+ \longrightarrow \mathcal{R}_K^+. \quad (72)$$

Lemma 27. *The trace map is characterized as follows:-*

$$\mathrm{Tr}_\phi^{(n)}(c \cdot \bigwedge_i d\log(T_i) \wedge d\log(\pi_K + 1)) = \frac{1}{p^n} \mathrm{Tr}_\phi(p.c\omega) \bigwedge_i d\log(T_i) \wedge d\log(\pi_K + 1), \quad (73)$$

where

$$\omega = \frac{\phi(\pi_K) + 1}{\pi_K + 1} \left(\frac{d}{d\pi_K} \phi(\pi_K + 1) \right)^{-1}. \quad (74)$$

Proof. First note that since ϕ is a ring homomorphism, we have $\frac{d}{d\pi_K}\phi(\pi_K+1) = \frac{d}{d\pi_K}\phi(\pi_K)$. Now by Corollary 13, there exists $I < 0$ and $a \in (\mathbb{A}_K^{\dagger,N})^\times$ for $N \gg 0$ such that

$$\frac{d}{d\pi_K}\phi(\pi_K) = p\pi_K^I a,$$

so $p\omega \in \mathcal{S}_K^{\dagger,N}$ for $N \gg 0$. The lemma is now an immediate consequence of Remark (iii) in Section 2.1 in [7]. $\square$

Note that $d\log(\pi_K) = \frac{\pi_K+1}{\pi_K}d\log(\pi_K+1)$. Using the formulae in Lemma 27, we can therefore extend the trace map to $\Omega_{\mathcal{R}_K^+}^n(\log)$:-

$$\mathrm{Tr}_\phi^{(n)}\left(\frac{c}{\pi_K} \cdot \bigwedge_i d\log(T_i) \wedge d\log(\pi_K+1)\right) = \frac{1}{p^n} \frac{\mathrm{Tr}_\phi(p \cdot c\tilde{\omega})}{\pi_K} \bigwedge_i d\log(T_i) \wedge d\log(\pi_K+1), \quad (75)$$

where

$$\tilde{\omega} = \frac{\phi(\pi_K)}{\pi_K} \frac{\phi(\pi_K)+1}{\pi_K+1} \left(\frac{d}{d\pi_K}\phi(\pi_K+1)\right)^{-1}. \quad (76)$$

Definition. Let $\psi : \Omega_{\mathcal{R}_K^+}^n(\log) \longrightarrow \Omega_{\mathcal{R}_K^+}^n(\log)$ be the map $\mathrm{Tr}_\phi^{(n)}$.

4.3 The logarithmic derivative

Since $\mathcal{R}_K^+$ is p -adically complete, the $d\log$ -operator induces a continuous map

$$d\log : \tilde{K}_n(\mathcal{S}_K) \longrightarrow \Omega_{\mathcal{R}_K^+}^n(\log) \quad (77)$$

which clearly factors through $\mathcal{V}$, giving

$$d\log : \tilde{K}_n(\mathcal{S}_K)/\mathcal{V} \longrightarrow \Omega_{\mathcal{R}_K^+}^n(\log). \quad (78)$$

Lemma 28. *We have a commutative diagram*

$$\begin{array}{ccc} \tilde{K}_n(\mathcal{S}_K)/\mathcal{V} & \xrightarrow{N} & \tilde{K}_n(\mathcal{S}_K)/\mathcal{V} \\ \downarrow d\log & & \downarrow d\log \\ \Omega_{\mathcal{R}_K^+}^n(\log) & \xrightarrow{\psi} & \Omega_{\mathcal{R}_K^+}^n(\log) \end{array}$$

Proof. Explicit computation. $\square$

Corollary 29. *We have a canonical homomorphism*

$$\tau : \tilde{K}_n(\mathbb{E}_K) \longrightarrow (\Omega_{\mathcal{R}_K^+}^n(\log))^{\psi=1}. \quad (79)$$

Proof. Define τ to be the composition of $d\log$ with the inverse of the isomorphism $(\tilde{K}_n(\mathcal{S}_K)/\mathcal{V})^{N=1} \cong \tilde{K}_n(\mathbb{E}_K)$. $\square$

To define the logarithmic derivative, we need to show the following result:-

Lemma 30. *The map τ extends to a map*

$$\tau : \widehat{K}_n(\mathbb{E}_{\mathcal{K}}) \longrightarrow (\Omega_{\mathcal{R}_K^+}^n(\log))^{\psi=1}. \quad (80)$$

Proof. Let $a \in \widehat{K}_n(\mathbb{E}_{\mathcal{K}})$. It follows from the construction of τ that $\tau(a)$ can be determined by the following procedure:- Let α be some lift of a to $\tilde{K}_n(\mathcal{S}_K)$. Then $\psi^k(d \log(\alpha)) \rightarrow \tau(a)$ as $k \rightarrow +\infty$. (One can show that $\psi^{k+1}(d \log(\alpha)) - \psi^k(d \log(\alpha)) \in p^k \Omega_{\mathcal{R}_K^+}^n(\log)$ for all k .)

For $r \geq 1$, let $\mathcal{U}^{(r)}$ be the subgroup of $\tilde{K}_n(\mathbb{E}_{\mathcal{K}})$ topologically generated by the symbols $\{a_1, \dots, a_n\}$ such that $a_1 \in 1 + \pi_K^r \mathbb{E}_{\mathcal{K}}^+$. Note that

$$\widehat{K}_n(\mathbb{E}_{\mathcal{K}}) = \varprojlim \tilde{K}_n(\mathbb{E}_{\mathcal{K}}) / \mathcal{U}^{(r)}. \quad (81)$$

Suppose that $a \in \ker(\tilde{K}_n(\mathbb{E}_{\mathcal{K}}) \rightarrow \widehat{K}_n(\mathbb{E}_{\mathcal{K}}))$, i.e. $a \in \bigcap_r \mathcal{U}^{(r)}$. Using the above algorithm for determining $\tau(a)$ then shows that $\tau(a) \in \bigcap_m p^m \Omega_{\mathcal{R}_K^+}^n(\log)$ and hence $\tau(a) = 0$ since $\Omega_{\mathcal{R}_K^+}^n(\log) \cong \mathcal{R}_K^+$. Now let $x \in \widehat{K}_n(\mathbb{E}_{\mathcal{K}})$, and write x as an (infinite) product $\prod_r x_r$, where $x_r \in \mathcal{U}^{(r)}$. For each r , choose a lift $\mathcal{X}_r$ in the subgroup of $\tilde{K}_n(\mathcal{S}_K)$ topologically generated by symbols of the form $\{b_1, \dots, b_n\}$ with $b_1 \in 1 + \pi_K^r O_H[[\pi_K]]$. Let $y_r = d \log(\mathcal{X}_r)$. Then $y_r \in \pi_K^r O_H[[\pi_K]] \wedge d \log(T_i) \wedge d \log(\pi_K)$, so the sum $\sum_r y_r$ converges to an element $y \in \Omega_{\mathcal{R}_K^+}^n(\log)$. Now the map Tr_{ϕ} is (p, π_K) -continuous, so for all k we have

$$\psi^k(y) = \psi^k(y_1) + \psi^k(y_2) + \dots \quad (82)$$

But we have $\psi^{k+1}(y_i) - \psi^k(y_i) \in p^k \Omega_{\mathcal{R}_K^+}^n(\log)$ for all i . Now $\Omega_{\mathcal{R}_K^+}^n(\log)$ is p -adically complete, so $\psi^k(y)$ converges towards some element $z \in \Omega_{\mathcal{R}_K^+}^n(\log)$ as $k \rightarrow +\infty$. Define $\tau(x) = z$. $\square$

Proposition 31. *We have a canonical homomorphism*

$$\varprojlim \widehat{K}_n(\mathcal{K}_m) \cong \widehat{K}_n(\mathbb{E}_{\mathcal{K}}). \quad (83)$$

Proof. For $i \geq N$, define the evaluation map

$$h_i^{(N)} : \mathcal{T}_K^{\dagger, N} \longrightarrow \mathcal{K}_i \quad (84)$$

by $h_i^{(N)}(\pi_K) = \pi_{K_n}$ and $h_i^{(N)}(T_j) = T_j^{\frac{1}{p^i}}$. Then for all $i \gg 0$, the map $h_i^{(N)}$ induces a homomorphism

$$H_i^{(N)} : \widehat{K}_n(\mathcal{T}_K^{\dagger, N}) \longrightarrow \widehat{K}_n(\mathcal{K}_i). \quad (85)$$

One can show that there exists $j_i^{(n)} \geq i - n - 1$ such that the image of $\ker(H_i^{(N)})$ in $\widehat{K}_n(\mathbb{E}_{\mathcal{K}})$ is the subgroup $\mathfrak{A}_i^{(N)}$ generated by the symbols of the

form $\{a_1, \dots, a_n\}$ with $a_1 \in 1 + \pi_K^{j_i^{(n)}} \mathbb{E}_K^+$, and we have commutative diagram

$$\begin{array}{ccccc}
K_n(\mathbb{E}_K)/\mathfrak{A}_{i+1}^{(N)} & \longleftarrow & \widehat{K}_n(\mathcal{T}_K^{\dagger, N})/\ker(H_{i+1}^{(N)}) & \xrightarrow[\cong]{H_{i+1}^{(N)}} & \widehat{K}_n(\mathcal{K}_{i+1}) \\
\downarrow & & \downarrow N & & \downarrow N_{O_{\mathcal{K}_{m+1}}/O_{\mathcal{K}_m}} \\
K_n(\mathbb{E}_K)/\mathfrak{A}_i^{(N)} & \longleftarrow & \widehat{K}_n(\mathcal{T}_K^{\dagger, N})/\ker(H_i^{(N)}) & \xrightarrow[\cong]{H_i^{(N)}} & \widehat{K}_n(\mathcal{K}_i)
\end{array}$$

Here, the map $K_n(\mathbb{E}_K)/\mathfrak{A}_{i+1}^{(N)} \rightarrow K_n(\mathbb{E}_K)/\mathfrak{A}_i^{(N)}$ is the natural projection map. Now $j_i^{(n)} \rightarrow +\infty$ as $i \rightarrow +\infty$, so we have $\varprojlim K_n(\mathbb{E}_K)/\mathfrak{A}_i^{(N)} = \widehat{K}_n(\mathbb{E}_K)$, and hence the above diagram induces a map

$$g : \varprojlim \widehat{K}_n(\mathcal{K}_i) \longrightarrow \widehat{K}_n(\mathbb{E}_K). \quad (86)$$

□

We can now define the logarithmic derivative:-

Definition. The logarithmic derivative is the map

$$\mathfrak{T} : \varprojlim \widehat{K}_n(O_{\mathcal{K}_m}) \longrightarrow (\Omega_{\mathcal{R}_K^+}^n(\log))^{\psi=1} \quad (87)$$

obtained by composing the homomorphism g with τ .

Problem. Recall that

$$\Omega_{\mathcal{R}_K^+}^n(\log) \cong \mathcal{R}_K^+ d. \log(\pi_K) \wedge d \log(T_1) \wedge \dots \wedge d \log(T_{n-1}). \quad (88)$$

In analogy to the construction of Cherbonnier and Colmez in the 1-dimensional case, we would like to associate to an element in $\varprojlim \widehat{K}_n(O_{\mathcal{K}_m})$ something like a function that is convergent on some annulus of the p -adic unit disc. However, the elements of $\mathcal{R}_K^+$ (as series in π_K) do not necessarily converge anywhere. Consequently, we need to define a suitable subring, an n -dimensional ring of overconvergent series. We will do that in the next section.

5 Higher dimensional rings of overconvergent series

5.1 Definition and properties

5.1.1 Definition of $\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}$

As in the classical case, we define the elements of $\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}$ as certain functions defined on some annulus on the p -adic open unit disc:- Let $x \in \mathcal{R}_{\mathbb{Q}_p}$, and write $x = \sum_{i \in \mathbb{Z}} f_i(T_1, \dots, T_{n-1}) \pi^i$.

Definition. Let $n \geq 1$. Then $x \in \mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}$ if the series $\sum_{i \in \mathbb{Z}} f_i(T_1, \dots, T_{n-1})\pi^i$ converges on the annulus $\{p^{-\frac{1}{(p-1)p^{N-1}}} \leq |X| < 1\} \subset \mathbb{C}_p$ (with values in $\mathbb{C}_p \hat{\otimes}_{\mathbb{Z}_p} O_{\mathcal{H}}$) and is bounded above by 1.

Alternatively, the elements of $\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}$ can be defined using growth conditions on the negative coefficients:-

Proposition 32. Let $x \in \mathcal{R}_{\mathbb{Q}_p}$, and write $x = \sum_{i \in \mathbb{Z}} f_i(T_1, \dots, T_{n-1})\pi^i$. Then $x \in \mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}$ if and only if $v(f_i) + \frac{i}{(p-1)p^{N-1}} \geq 0$ for all $i < 0$ and $i \rightarrow +\infty$ as $i \rightarrow -\infty$.

Proof. Explicit calculation. □

Corollary 33. The ring $\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}$ is an $\mathbb{A}_{\mathbb{Q}_p}^{\dagger, N}$ -module.

It follows from the definition and the properties of the ring $\mathbb{A}_{\mathbb{Q}_p}^{\dagger, N}$ that we have the following corollary:-

Corollary 34. (1) $\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N} \subset \mathcal{R}_{\mathbb{Q}_p}^{\dagger, N+1}$, and
(2) If $x \in \mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}$, then $\phi(x) \in \mathcal{R}_{\mathbb{Q}_p}^{\dagger, N+1}$.

Proof. To prove (2), observe that if $\alpha \in \{p^{-\frac{1}{(p-1)p^N}} \leq |X| < 1\}$, then $(\alpha+1)^p - 1 \in \{p^{-\frac{1}{(p-1)p^{N-1}}} \leq |X| < 1\}$. □

Proposition 35. The ring $\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N+1}$ is a free $\phi(\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N})$ -module of degree p^n .

Proof. Explicit calculation shows that $\{T_1^{i_1} \dots T_{n-1}^{i_{n-1}}(\pi+1)^j\}_{0 \leq i_1, \dots, i_{n-1}, j \leq p-1}$ is a basis of $\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N+1}$ over $\phi(\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N})$. □

Definition. The ring $\mathcal{R}_{\mathbb{Q}_p}^{\dagger} \subset \mathcal{R}_{\mathbb{Q}_p}$ of overconvergent elements is the union

$$\mathcal{R}_{\mathbb{Q}_p}^{\dagger} = \bigcup \mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}.$$

The topology on $\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}$ is induced from the topology on $\mathcal{R}_{\mathbb{Q}_p}$:-

Lemma 36. A set of neighbourhoods of 0 in $\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}$ is given by $p^k \mathcal{R}_{\mathbb{Q}_p}^{\dagger, N} + \pi^l \mathcal{R}_{\mathbb{Q}_p}^{\dagger}$.

Remark. The topology on $\mathcal{R}_{\mathbb{Q}_p}^{\dagger}$ is the Frechet topology induced from the topologies on the $\mathcal{R}_{\mathbb{Q}_p}^{\dagger, N}$.

5.1.2 Definition of $\mathcal{R}_K^{\dagger,N}$

Since $\mathbb{A}_K^{\dagger,N}$ is a free finitely generated module over $\mathbb{A}_{\mathbb{Q}_p}^{\dagger,N}$ for $N \gg 0$, it seems natural to make the following definition:-

Definition. $\mathcal{R}_K^{\dagger,N} = \mathbb{A}_K^{\dagger,N} \otimes_{\mathbb{A}_{\mathbb{Q}_p}^{\dagger,N}} \mathcal{R}_{\mathbb{Q}_p}^{\dagger,N}$

Lemma 37. (1) $\mathcal{R}_K^{\dagger,N} \subset \mathcal{R}_K^{\dagger,N+1}$,
(2) if $x \in \mathcal{R}_K^{\dagger,N}$, then $\phi(x) \in \mathcal{R}_K^{\dagger,N+1}$, and
(3) $\mathcal{R}_K^{\dagger,N+1}$ is a free finitely generated module of degree p^n over $\phi(\mathcal{R}_K^{\dagger,N})$.

Proof. (1) and (2) are immediate consequences of Corollary 34 and the properties of the ring $\mathbb{A}_K^{\dagger,N}$. Now as shown in [3], Lemma V.1.4, the set $\{1, \dots, (\pi + 1)^{p-1}\}$ is a basis of $\mathbb{A}_K^{\dagger,N+1}$ over $\phi(\mathbb{A}_K^{\dagger,N})$. Together with Proposition 35, this proves (3). $\square$

Let $\mathcal{R}_K^+ = \mathbb{A}_K^+ \otimes_{\mathbb{A}_{\mathbb{Q}_p}^+} \mathcal{R}_{\mathbb{Q}_p}^+$.

Lemma 38. A set of neighbourhoods of 0 in $\mathcal{R}_K^{\dagger,N}$ is given by $p^k \mathcal{R}_K^{\dagger,N} + \pi^l \mathcal{R}_K^+$ for $l, k \geq 1$.

If $n \gg 0$, then the elements of $\mathcal{R}_K^{\dagger,N}$ also have a description as functions defined on some annulus on the p -adic open unit disc:- Now let $x \in \mathcal{R}_K^{\dagger,N}$, and let $n(K)$ be as defined in Proposition 6. We can then write x as $\sum_{i \in \mathbb{Z}} f_i(T_1, \dots, T_{n-1}) \pi_K^i$.

Proposition 39. Let $N \geq n(K)$. Then $x \in \mathcal{R}_K^{\dagger,N}$ if and only if the series $\sum_{i \in \mathbb{Z}} f_i(T_1, \dots, T_{n-1}) X^i$ converges on the annulus $\{p^{-\frac{1}{e(p-1)p^{N-1}}} \leq |X| < 1\}$ and is bounded above by 1.

To prove the proposition, we need the following result:-

Lemma 40. We have that $x \in \mathcal{R}_K^{\dagger,N}$ if and only if $v(f_i) + \frac{i}{e(p-1)p^{N-1}} \geq 0$ for all $i < 0$ and $\rightarrow +\infty$ as $i \rightarrow -\infty$.

Proof. Imitate the proof of Proposition 32. $\square$

Proof. Note that $1, \pi_K, \dots, \pi_K^{e-1}$ is a basis for $\mathbb{A}_K^{\dagger,N}$ over $\mathbb{A}_{\mathbb{Q}_p}^{\dagger,N}$. Recall from (40) that there exist $c_i \in \mathbb{Z}_p$ such that

$$\pi_K^e = \pi(c_0 + c_1 \pi_K + c_2 \pi_K^2 + \dots). \quad (89)$$

It follows that if we write

$$\pi_K^e = a_0 + a_1 \pi_K + \dots + a_{e-1} \pi_K^{e-1} \quad (90)$$

with $a_i \in \mathbb{A}_{\mathbb{Q}_p}^{\dagger,N}$, then $a_i \bmod p \in \bar{\pi}_K \mathbb{E}_{\mathbb{Q}_p}^+$, i.e. as a function on $\{p^{-\frac{1}{(p-1)p^{N-1}}} \leq |X| < 1\}$, a_i is strictly less than 1 for all i . Now the topology on $\mathbb{A}_{\mathbb{Q}_p}^{\dagger,N}$ agrees

with the topology of uniform convergence on every compact subannulus of $\{p^{-\frac{1}{(p-1)p^{N-1}}} \leq |X| < 1\}$. It follows that if we write

$$\pi_K^j = a_{j,0} + a_{j,1}\pi_K + \cdots + a_{j,e-1}\pi_K^{e-1} \quad (91)$$

with $a_{j,k} \in \mathbb{A}_{\mathbb{Q}_p}^{\dagger,N}$, then $a_{j,k} \rightarrow 0$ as $j \rightarrow +\infty$ for all $0 \leq k < e$.

Suppose now that $\sum_{j \in \mathbb{Z}} f_j(T_1, \dots, T_{n-1})X^j$ converges on $\{p^{-\frac{1}{e(p-1)p^{N-1}}} \leq |X| < 1\}$ and is bounded above by 1. Write

$$\sum_{j \in \mathbb{Z}} f_j(T_1, \dots, T_{n-1})X^j = \sum_{i=0}^{e-1} X^i \sum_{j \in \mathbb{Z}} f_j(T-1, \dots, T_{n-1})a_{j,i}. \quad (92)$$

Now $v(f_j) > 0$ for all $j < 0$ and $\rightarrow +\infty$ as $j \rightarrow -\infty$ by the previous lemma, so $f_j(T_1, \dots, T_{n-1})a_{j,i}$ is strictly less than 1 on $\{p^{-\frac{1}{(p-1)p^{N-1}}} \leq |X| < 1\}$ and $\rightarrow 0$ as $j \rightarrow -\infty$. Since we have shown above that $a_{j,k} \rightarrow 0$ as $j \rightarrow +\infty$, it follows that the series $\sum_{j \in \mathbb{Z}} f_j(T-1, \dots, T_{n-1})a_{j,i}$ converges on $\{p^{-\frac{1}{(p-1)p^{N-1}}} \leq |X| < 1\}$ and is bounded above by 1 and hence is an element of $\mathcal{R}_{\mathbb{Q}_p}^{\dagger,N}$.

Conversely, if $x \in \mathcal{R}_K^{\dagger,N}$, then it follows from the definition of $\mathcal{R}_K^{\dagger,N}$ and Proposition 6 that the expansion of x as a series in π_K converges on the annulus $\{p^{-\frac{1}{e(p-1)p^{N-1}}} \leq |X| < 1\}$ and is bounded above by 1. $\square$

Definition. Define the ring of overconvergent elements of $\mathcal{R}_K$ as

$$\mathcal{R}_K^{\dagger} = \bigcup_{n \geq n(K)} \mathcal{R}_K^{\dagger,N}.$$

5.1.3 Properties of $\mathcal{R}_K^{\dagger}$

Throughout this section, we assume that $N \geq n(K)$.

Definition. In the (p, π_K) -adic topology on $\mathcal{R}_K^{\dagger,N}$, a basis of neighbourhoods of 0 is given by $p^k \mathcal{R}_K^{\dagger,N} + \pi_K^l O_K[[\pi_K]]$, where $k, l \geq 1$.

Lemma 41. *The (p, π) -adic topology and the (p, π_K) -adic topology agree on $\mathcal{R}_K^{\dagger,N}$.*

Proof. Recall that $\mathcal{R}_K^{\dagger,N} = \mathbb{A}_K^{\dagger,N} \otimes_{\mathbb{A}_{\mathbb{Q}_p}^{\dagger,N}} \mathcal{R}_{\mathbb{Q}_p}^{\dagger,N}$. The lemma is now a consequence of the fact that the (p, π) -adic topology and the (p, π_K) -adic topology agree on $\mathbb{A}_K^{\dagger,N}$ (c.f. Section 2.6 in [1]). $\square$

Remark. The (p, π_K) -adic topology is the topology of convergence on compact subannuli of $\{p^{-\frac{1}{e(p-1)p^{N-1}}} \leq |X| < 1\}$.

Lemma 42. *The ring $\mathcal{R}_K^{\dagger,N}$ is complete with respect to the (p, π_K) -adic topology.*

Proof. Clear from Proposition 39 (or the remark above). $\square$

Lemma 43. *Let $x \in \mathcal{R}_K^{\dagger, N}$ and write $x = \sum f_i(T_1, \dots, T_{n-1})\pi_K^i$. If $v(f_0) = 0$ and $v(f_i) + \frac{i}{(p-1)p^{N-1}} > 0$ for all $i < 0$, then x is invertible in $\mathcal{R}_K^{\dagger, N}$.*

Proof. Suppose that $v(f_0) = 0$ and $v(f_i) + \frac{i}{e(p-1)p^{N-1}} > 0$ for all $i > 0$. Since $O_{\mathcal{H}}$ is a p -adically complete local ring, f_0 is invertible in $O_{\mathcal{H}}$ (so wlog = 1) and hence $\sum_{i \geq 0} f_i(T)\pi_K^i$ is invertible in $\mathcal{R}_K^{\dagger, N}$ (since the formal inverse converges), so without loss of generality assume that $x = 1 + \sum_{j \leq -1} f_j(T_1, \dots, T_{n-1})\pi_K^j$. The formal inverse of x is given by $\sum_{k \geq 0} (1-x)^k = \sum_{i \in \mathbb{Z}} h_i(T_1, \dots, T_{n-1})\pi_K^i$. We need to show that this sum converges in $\mathcal{R}_K^{\dagger, N}$. Since $v(f_j) + \frac{j}{e(p-1)p^{N-1}} > 0$ for all j , it is clear from the formal expansion that $v(h_j) + \frac{j}{e(p-1)p^{N-1}} > 0$ for all $j < 0$. It remains to show that $v(h_j) + \frac{j}{e(p-1)p^{N-1}} \rightarrow +\infty$ as $j \rightarrow -\infty$. Let $\alpha = \frac{1}{e(p-1)p^{N-1}}$. Since $v(f_i)$ is a positive integer for all i , $v(f_i) + \frac{i}{e(p-1)p^{N-1}} \geq \alpha$ for all $i > 0$. Fix $M > 0$, and let $K \gg 0$ such that $v(f_i) + \frac{i}{e(p-1)p^{N-1}} \geq M\alpha$ for all $i \geq K$. Expanding $\sum_{k \geq 0} (1-x)^k$ then shows that $v(h_j) + \frac{j}{e(p-1)p^{N-1}} \geq M\alpha$ for all $j \geq MN$, which finishes the proof. $\square$

Lemma 44. *Let $N \geq n(K)$. Let $x = \sum_{i \geq 0} f_i(T_1, \dots, T_{n-1})\pi_K^i \in \mathcal{R}_K^{\dagger, N}$. Then the following are equivalent:-*

- (i) x is invertible in $\mathcal{R}_K^{\dagger}$,
- (ii) x is invertible in $\mathcal{R}_K^{\dagger, N+1}$,
- (iii) $v(f_0) = 0$.

Proof. It follows from Lemma 43 that if $v(f_0) = 0$, then x is invertible in $\mathcal{R}_K^{\dagger, N+1}$. Conversely, write $x = \sum_{i \in \mathbb{Z}} f_i(T_1, \dots, T_{n-1})\pi_K^i$, and suppose that x is invertible in $\mathcal{R}_K^{\dagger}$. Since $p^{-1} \notin \mathcal{R}_K$, we must have $v(f_0) = 0$, i.e. f_0 is a unit in $O_{\mathcal{H}}$ and hence invertible in $\mathcal{R}_K^{\dagger, N}$. $\square$

Lemma 44 has the following important consequence:-

Proposition 45. $\mathcal{R}_K^{\dagger}$ can be written as the (increasing) union of p -adically complete local rings.

Proof. Let $N \geq n(K)$. To simplify the notation, write (only in this proof) R_N for $\mathcal{R}_K^{\dagger, N}$. Recall that every element of R_N can be written as a power series $\sum_{i \in \mathbb{Z}} f_i(T_1, \dots, T_{n-1})\pi_K^i$. Let

$$\mathfrak{p}_N = \left\{ \sum_{i \in \mathbb{Z}} f_i(T_1, \dots, T_{n-1})\pi_K^i \in R_N \mid v(f_0) > 0 \right\}.$$

Then $\mathfrak{p}_N$ is an ideal of R_N , and explicit calculation shows that it is prime. Note that $p \in \mathfrak{p}_N$. Let $R_{N, \mathfrak{p}_N}$ denote the localisation of R_N at $\mathfrak{p}_N$, which by Lemma 44 is contained in R_{N+1} , and let S_N be its completion with respect to $\mathfrak{p}_N$. Then $\mathfrak{p}_N$ is still a local ring, and it is p -adically complete since $p \in \mathfrak{p}_N$. The

topology defined by $\mathfrak{p}_N$ is the (p, π_K) -adic topology. To finish the proof of the proposition, note that $S_N \subset R_{N+1}$ since R_{N+1} is complete in the (p, π_K) -adic topology (and hence the topology on S_N is the subspace topology induced from R_{N+1}), i.e. $\mathcal{R}_K^\dagger = \bigcup_{N \geq n(K)} S_N$. $\square$

Definition. Write $\mathcal{S}_K^{\dagger, N}$ for the ring S_N .

Definition. Let $\mathcal{T}_K^{\dagger, N} = \mathcal{S}_K^{\dagger, N}[\pi_K^{-1}]$.

Note. (1) Note that $\mathcal{T}_K^{\dagger, N} = \mathcal{S}_K^{\dagger, N}[\pi^{-1}]$.

(2) Written as power series in π_K , the elements of $\mathcal{T}_K^{\dagger, N}$ can be interpreted as functions converging on $\{p^{-\frac{1}{(p-1)\epsilon_K p^{N-1}}} \leq |X| < 1\}$ (but not necessarily bounded above by 1).

Remark. Since Frobenius of the residue field is the p -power map, we have $\phi(\mathcal{S}_K^{\dagger, N}) \subset \mathcal{S}_K^{\dagger, N+1}$ and $\phi(\mathcal{T}_K^{\dagger, N}) \subset \mathcal{T}_K^{\dagger, N+1}$.

5.2 Properties of ϕ

Proposition 46. Let $N \gg 0$. Then $\mathcal{R}_K^{\dagger, N+1}$ is a free $\phi(\mathcal{R}_K^{\dagger, N})$ -module of rank p^N .

Proof. It follows from the classical theory of the rings $\mathbb{A}_{\mathbb{Q}_p}^{\dagger, N}$ and $\mathbb{A}_K^{\dagger, N}$ that $\{T_1^{i_1} \dots T_{n-1}^{i_{n-1}} (1 + \pi)^j\}_{0 \leq i_1, \dots, i_{n-1}, j \leq p-1}$ is a basis of $\mathcal{R}_K^{\dagger, N+1}$ over $\phi(\mathcal{R}_K^{\dagger, N})$. $\square$

Corollary 47. $\mathcal{R}_K^\dagger$ is a free $\phi(\mathcal{R}_K^\dagger)$ -module of rank p^n .

The following proposition is the reason why the ring of overconvergent series can be used to construct the logarithmic derivative:-

Proposition 48. For $n \gg 0$, $\mathcal{T}_K^{\dagger, N+1}$ is a free $\phi(\mathcal{T}_K^{\dagger, N})$ -module of rank p^n .

Noting that $\mathcal{T}_K^{\dagger, N} = \mathcal{S}_K^{\dagger, N}[\pi^{-1}]$, this proposition is a consequence of the following result:-

Proposition 49. For $N \gg 0$, $\mathcal{S}_K^{\dagger, N+1}$ is a free $\phi(\mathcal{S}_K^{\dagger, N})$ -module of rank p^n .

To prove the Proposition 49, we begin with the following observations:-

Lemma 50. Let $N \in \mathbb{N}$. If we write $\pi^N = \sum_{0 \leq i \leq p-1} \phi(a_i^{(N)})(1 + \pi)^i$, with $a_i^{(N)} \in \mathbb{A}_{\mathbb{Q}_p}^+$, then for all $0 \leq i \leq p-1$ we have $a_i^{(N)} \rightarrow 0$ as $N \rightarrow +\infty$.

Proof. $a_i^{(N)} = \psi(\pi^N(1 + \pi)^{-i})$, and ψ is continuous on $\mathbb{A}_{\mathbb{Q}_p}$. $\square$

Remark. It follows that if we expand π^N as $\sum_{0 \leq i \leq p-1} \phi(b_i^{(N)})\pi^i$, then $b_i^{(N)} \rightarrow 0$ as $N \rightarrow +\infty$.

We also need the following lemma:-

Lemma 51. $\pi^i \rightarrow 0$ in $\mathcal{S}_K^{\dagger, N}$ as $i \rightarrow +\infty$ for all $N \gg 0$.

Proof. Recall that $\pi = \pi_K^e(b_0 + b_1\pi_K + b_2\pi_K^2 + \dots)$ for some $b_i \in \mathbb{Z}_p$. $\square$

We can now prove Proposition 49:-

Proof. Wlog assume that ϕ acts trivially on $T_1, \dots, T_{n-1}$, so $\{1, \pi, \dots, \pi^{p-1}\}$ is a basis for $\mathcal{R}_K^{\dagger, N+1}$ over $\mathcal{R}_K^{\dagger, N}$. We first show that every element of $(\mathcal{R}_K^{\dagger, n+1})_{\mathfrak{p}_{n+1}}$ can be written as $\sum_{0 \leq i \leq p-1} \phi(x_i)\pi^i$ with $x_i \in \mathcal{S}_K^{\dagger, N}$.

Let $x \in \mathcal{R}_K^{\dagger, N+1}$ and write $x = \sum_{i \in \mathbb{Z}} f_i(T_1, \dots, T_{n-1})\pi_K^i$. Assume that $v(f_0) = 0$. Let $x_0, \dots, x_{p-1} \in \mathcal{R}_K^{\dagger, N}$ such that $x = \sum_{0 \leq j < p} \phi(x_j)\pi^j$. Since $v(f_0) = 0$, $x_0^{-1} \in (\mathcal{R}_K^{\dagger, N})_{\mathfrak{p}_n}$, so $\phi(x_0^{-1}) \in (\mathcal{R}_K^{\dagger, N+1})_{\mathfrak{p}_{N+1}}$. We can therefore write $x = \phi(x_0)(1 + \pi\phi(x_0^{-1})z)$, where $z = \sum_{0 < i \leq p-1} \phi(x_i)\pi^{i-1}$. Let

$$y = 1 - \pi\phi(x_0^{-1})z + \pi^2\phi(x_0^{-2})z^2 - \pi^3\phi(x_0^{-3})z^3 + \dots \quad (93)$$

be the formal inverse of $1 + \pi\phi(x_0^{-1})z$. Since $1 + \pi\phi(x_0^{-1})z \in \mathcal{R}_K^{\dagger, N+1} \setminus \mathfrak{p}_{N+1}$, we have $y \in \mathcal{R}_K^{\dagger, N+2}$, so there exist $y_0, \dots, y_{p-1} \in \mathcal{R}_K^{\dagger, N+1}$ such that $y = \sum_{0 \leq k \leq p-1} \phi(y_k)\pi^k$. We will show that in fact $y_k \in \mathcal{S}_K^{\dagger, N}$ for all k . Let $0 \leq k \leq p-1$. By expanding (93) (and remembering that $\phi(\pi) = (\pi+1)^p - 1$), we see that y_k is of the form

$$y_k = \sum_{l,m=0}^{\infty} p^l \pi^m \alpha_{l,m}^{(k)} \quad (94)$$

for some $\alpha_{l,m}^{(k)} \in (\mathcal{R}_K^{\dagger, N})_{\mathfrak{p}_N}$. But $p^l \pi^m \rightarrow 0$ in $\mathcal{S}_K^{\dagger, N}$ as $l, m \rightarrow +\infty$ by Lemma 51, so since $\mathcal{S}_K^{\dagger, N}$ is complete, it follows that $y_k \in \mathcal{S}_K^{\dagger, N}$. Since ϕ is a ring homomorphism, we have shown that every element $z \in (\mathcal{R}_K^{\dagger, N+1})_{\mathfrak{p}_{N+1}}$ can be written as $\sum_{0 \leq i \leq p-1} \phi(z_i)\pi^i$ with $z_i \in \mathcal{S}_K^{\dagger, N}$.

It remains to show that this property extends to $\mathcal{S}_K^{\dagger, N+1}$. Identify $\mathfrak{p}_{N+1}$ with its image in $(\mathcal{R}_K^{\dagger, N+1})_{\mathfrak{p}_{N+1}}$, and assume that $x \in \mathfrak{p}_{N+1}^M$ for some $M \gg 0$. Write $\sum_{0 \leq i \leq p-1} \phi(x_i)\pi^i$ with $x_i \in \mathcal{S}_K^{\dagger, N}$. By completeness, it is sufficient to show that $x_i \in \mathfrak{p}_n^L \mathcal{S}_K^{\dagger, N}$ for some $L \gg 0$. Now since $x \in \mathfrak{p}_{N+1}^M$, there exist $a_1 \in \mathcal{R}_K^{\dagger, N+2}$, $a_2 \in \mathcal{R}_K^+$ and $N_1, N_2 \gg 0$ such that $x = p^{N_1}a_1 + \pi^{N_2}a_2$. Since ϕ is $\mathbb{Z}_p$ -linear, we can wlog. assume that $a_1 = 0$, so $x = \pi^{N_2}a_2$. But if we write $\pi^{N_2} = \sum_{0 \leq i \leq p-1} \phi(b_i^{(N_2)})\pi^i$, then we deduce from Lemma 50 that there exists some $L \gg 0$ such that $b_i^{(N_2)} \in \mathfrak{p}_n^L$ for all i . Since $\mathfrak{p}_n^L$ is an ideal, it follows that $x_i \in \mathfrak{p}_n^L$ for all $0 \leq i \leq p-1$, which finishes the proof. $\square$

5.3 Modules of differentials

We make the following definitions:-

Definition. Let $\Omega_{\mathcal{S}_K^{\dagger, N}/\mathbb{Z}}^1$ be the module of absolute differential forms of $\mathcal{S}_K^{\dagger, N}$. For $r \geq 1$, let $\Omega_{\mathcal{S}_K^{\dagger, N}/\mathbb{Z}}^r = \bigwedge_{\mathcal{S}_K^{\dagger, N}/\mathbb{Z}}^r \Omega_{\mathcal{S}_K^{\dagger, N}/\mathbb{Z}}^1$, and define $\Omega_{\mathcal{S}_K^{\dagger, N}}^r$ to be its p -adic

completion

$$\Omega_{\mathcal{S}_K^{\dagger,N}}^r = \varprojlim \Omega_{\mathcal{S}_K^{\dagger,N}/\mathbb{Z}}^r / p^m \Omega_{\mathcal{S}_K^{\dagger,N}/\mathbb{Z}}^r. \quad (95)$$

Definition. Let

$$\Omega_{\mathcal{S}_K^{\dagger,N}/\mathbb{Z}}^1(\log) = \Omega_{\mathcal{S}_K^{\dagger,N}/\mathbb{Z}}^1 \oplus \mathcal{S}_K^{\dagger,N} \otimes_{\mathbb{Z}} (\mathcal{T}_K^{\dagger,N})^{\times} / \mathcal{N}, \quad (96)$$

where $\Omega_{\mathcal{S}_K^{\dagger,N}/\mathbb{Z}}^1$ is the module of absolute differential forms and $\mathcal{N}$ is the submodule of the direct sum generated by elements $(-da, a \otimes a)$ for $a \in \mathcal{S}_K^{\dagger,N} \cap (\mathcal{T}_K^{\dagger,N})^{\times}$. In $\Omega_{\mathcal{S}_K^{\dagger,N}/\mathbb{Z}}^1(\log)$, denote the class $(0, 1 \otimes a)$ by $d \log a$. For $r \geq 1$, let $\Omega_{\mathcal{S}_K^{\dagger,N}/\mathbb{Z}}^r(\log) = \bigwedge_{\mathcal{S}_K^{\dagger,N}}^r \Omega_{\mathcal{S}_K^{\dagger,N}/\mathbb{Z}}^1(\log)$, and define $\Omega_{\mathcal{S}_K^{\dagger,N}}^r(\log)$ to be its p -adic completion

$$\Omega_{\mathcal{S}_K^{\dagger,N}}^r(\log) = \varprojlim \Omega_{\mathcal{S}_K^{\dagger,N}/\mathbb{Z}}^r(\log) / p^m \Omega_{\mathcal{S}_K^{\dagger,N}/\mathbb{Z}}^r(\log). \quad (97)$$

Lemma 52. *We have the following isomorphisms of $\mathcal{S}_K^{\dagger,N}$ -modules:-*

$$\Omega_{\mathcal{S}_K^{\dagger,N}}^n \cong \mathcal{S}_K^{\dagger,N} \cdot \bigwedge d \log(T_i) \wedge d \log(\pi_K + 1), \quad (98)$$

$$\Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log) \cong \mathcal{S}_K^{\dagger,N} \cdot \bigwedge d \log(T_i) \wedge d \log(\pi_K). \quad (99)$$

Proof. Follows from explicit computation, using Lemma 10 and the observation that $\pi_K + 1$ is invertible in $\mathcal{S}_K^{\dagger,N}$. $\square$

Now $\mathcal{S}_K^{\dagger,n+1}$ is finite flat and locally of complete intersection over $\phi(\mathcal{S}_K^{\dagger,N})$, so as shown in [7], we have a trace map of differential forms

$$\mathrm{Tr}_{\phi}^{(n)} : \Omega_{\mathcal{S}_K^{\dagger,N+1}}^n \longrightarrow \Omega_{\mathcal{S}_K^{\dagger,N}}^n. \quad (100)$$

and the usual trace map

$$\mathrm{Tr}_{\phi} : \mathcal{S}_K^{\dagger,N+1} \longrightarrow \mathcal{S}_K^{\dagger,N}. \quad (101)$$

Lemma 53. *The trace map is characterized as follows:-*

$$\mathrm{Tr}_{\phi}^{(n)}(c \cdot \bigwedge d \log(T_i) \wedge d \log(\pi_K + 1)) = \frac{1}{p^2} \mathrm{Tr}_{\phi}(p \cdot c \omega) \bigwedge d \log(T_i) \wedge d \log(\pi_K + 1), \quad (102)$$

where

$$\omega = \frac{\phi(\pi_K) + 1}{\pi_K + 1} \left(\frac{d}{d\pi_K} \phi(\pi_K + 1) \right)^{-1}. \quad (103)$$

Proof. First note that since ϕ is a ring homomorphism, we have $\frac{d}{d\pi_K} \phi(\pi_K + 1) = \frac{d}{d\pi_K} \phi(\pi_K)$. Now by Corollary 13, there exists $I < 0$ and $a \in (\mathbb{A}_K^{\dagger,M})^{\times}$ for $M \gg 0$ such that

$$\frac{d}{d\pi_K} \phi(\pi_K) = p \pi_K^I a,$$

so $p\omega \in \mathcal{S}_K^{\dagger,M}$ for $M \gg 0$. The lemma is now an immediate consequence of Remark (iii) in Section 2.1 in [7]. $\square$

Using these formulae, we extend the trace map to $\Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log)$:-

$$\mathrm{Tr}_\phi^{(n)}\left(\frac{c}{\pi_K} \cdot \bigwedge d\log(T_i) \wedge d\log(\pi_K+1)\right) = \frac{1}{p^n} \frac{\mathrm{Tr}_\phi(p \cdot c\tilde{\omega})}{\pi_K} \bigwedge d\log(T_i) \wedge d\log(\pi_K+1), \quad (104)$$

where

$$\tilde{\omega} = \frac{\phi(\pi_K)}{\pi_K} \frac{\phi(\pi_K)+1}{\pi_K+1} \left(\frac{d}{d\pi_K} \phi(\pi_K+1)\right)^{-1}. \quad (105)$$

Definition. Let $\psi : \Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log) \longrightarrow \Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log)$ be the map obtained by composing the natural map $\Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log) \longrightarrow \Omega_{\mathcal{S}_K^{\dagger,N+1}}^n(\log)$ with $\mathrm{Tr}_\phi^{(n)}$.

6 The logarithmic derivative revisited

Lemma 54. *The natural map $\iota : \Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log) \longrightarrow \Omega_{\mathcal{R}_K^+}^n(\log)$ is a closed embedding.*

Proof. Recall that

$$\Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log) \cong \mathcal{S}_K^{\dagger,N} d\log(\pi_K) \wedge d\log(T), \quad (106)$$

$$\Omega_{\mathcal{R}_K^+}^n(\log) \cong \mathcal{R}_K^+ d\log(\pi_K) \wedge d\log(T) \quad (107)$$

The map ι is just the natural inclusion $\mathcal{S}_K^{\dagger,N} \hookrightarrow \mathcal{R}_K^+$, which is a closed embedding. $\square$

As a corollary, we get the desired information about the logarithmic derivative:-

Theorem 55. *The logarithmic derivative gives a canonical homomorphism*

$$\mathfrak{T} : \varprojlim \widehat{K}_n(\mathcal{K}_i) \longrightarrow (\Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log))^{\psi=1}. \quad (108)$$

Proof. It is certainly sufficient to show that we get a canonical homomorphism

$$\widehat{K}_n(\mathbb{E}_{\mathcal{K}}) \longrightarrow (\Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log))^{\psi=1}. \quad (109)$$

Now if $\tilde{x} \in \widehat{K}_n(\mathbb{E}_{\mathcal{K}})$, then since $n \geq n(K)$ we can choose a lift x of $\tilde{x}$ such that x can be written as a product of symbols $\{b_1, \dots, b_n\}$ such that $b_i \in \mathcal{T}_K^{\dagger,N}$ for all i . Then $d\log(x) \in \Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log)$, and the sequence $(\psi^k(d\log(x)))$ then converges to $y = \tau(\tilde{x}) \in (\Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log))^{\psi=1}$. But ψ restricts to $\mathrm{Im}(\iota)$, so since $\Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log) \longrightarrow \Omega_{\mathcal{R}_K^+}^n(\log)$ is a closed embedding, it follows that $y \in \Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log)$, which finishes the proof. $\square$

We have the following special case when $K = \mathbb{Q}_p$:-

Corollary 56. *Suppose that $\mathcal{K} = \mathcal{H}$, i.e. $K = \mathbb{Q}_p$. Then the logarithmic derivative gives a canonical homomorphism*

$$\mathfrak{T} : \varprojlim \widehat{K}_n(\mathcal{H}_n) \longrightarrow (\Omega_{O_{\mathcal{H}}[[\pi]]}^n(\log))^{\psi=1}. \quad (110)$$

Remark. Note that

$$\Omega_{O_{\mathcal{H}}[[\pi]]}^n(\log) \cong O_{\mathcal{H}}[[\pi]] \cdot \bigwedge d \log(T_i) \wedge d \log(\pi). \quad (111)$$

7 Relation to the dual exponential map

7.1 Statement of the theorem

Let $N \geq n(K)$. In this section we will use the logarithmic derivative to give a new description of Kato's dual exponential map λ_m for $m \gg N$. Recall that for all $m \geq 1$, the homomorphism λ_m is defined as the composition

$$\lambda_m : \varprojlim \widehat{K}_n(\mathcal{K}_i) \xrightarrow{(\delta_{\mathcal{K}_i, p^i}^n)_{i \geq 1}} \varprojlim H^n(\mathcal{K}_i, (\mathbb{Z}/p^i\mathbb{Z})(n)) \quad (112)$$

$$\xrightarrow{\zeta_{p^i}^{\otimes(-1)}} \varprojlim H^n(\mathcal{K}_i, (\mathbb{Z}/p^i\mathbb{Z})(n-1)) \quad (113)$$

$$\xrightarrow{\text{trace}} \varprojlim H^n(\mathcal{K}_m, (\mathbb{Z}/p^i\mathbb{Z})(n-1)) \quad (114)$$

$$= H^n(\mathcal{K}_m, \mathbb{Z}_p(n-1)) \quad (115)$$

$$\longrightarrow H^n(\mathcal{K}_m, \mathbb{C}_p(n-1)) \cong \Omega_{\mathcal{K}_m}^{n-1}, \quad (116)$$

where $\delta_{\mathcal{K}_i, p^i}^n$ is the Galois symbol. Now let $\mathfrak{d}_m$ be the homomorphism

$$\mathfrak{d}_m : \varprojlim \widehat{K}_n(\mathcal{K}_i) \xrightarrow{\mathfrak{T}} (\Omega_{\mathcal{S}_K^{\dagger, N}}^n(\log))^{\psi=1} \quad (117)$$

$$\hookrightarrow \Omega_{\mathcal{S}_K^{\dagger, N}}^n(\log) \cong \frac{\mathcal{S}_K^{\dagger, N}}{\pi_K} \cdot \bigwedge d \log(T_i) \wedge d \log(\pi_K + 1) \quad (118)$$

$$\longrightarrow \Omega_{\mathcal{K}_m}^{n-1}. \quad (119)$$

Here the last arrow is defined by

$$f \cdot \bigwedge d \log(T_i) \wedge d \log(\pi_K + 1) \longrightarrow \frac{1}{p^m} h_m^{(N)}(f) \cdot \bigwedge d \log(h_m^{(N)}(T_i)) \quad (120)$$

where

$$h_m^{(N)} : \mathcal{S}_K^{\dagger, N} \rightarrow O_{\mathcal{K}_m} \quad (121)$$

is determined by $h_m^{(N)}(T_i) = T_i^{\frac{1}{p^m}}$ and $h_m^{(N)}(\pi_K) = \pi_{\mathcal{K}_m}$. (Note that this is well-defined for $m \geq N$ since the elements of $\mathcal{S}_K^{\dagger, N}$ can be interpreted as functions converging on the annulus $\{p^{-\frac{1}{e(p-1)p^N}} \leq |X| < 1\} \subset \mathbb{C}_p$.) We then have the following result:-

Theorem 57. *For all $m \geq N$, we have*

$$\mathfrak{d}_m = -\lambda_m. \quad (122)$$

7.2 Proof

7.2.1 Evaluation of the logarithmic derivative

Definition. For a complete discrete valuation field L of mixed characteristic $(0, p)$, let

$$\Omega_{O_L/\mathbb{Z}}^1(\log) = (\Omega_{O_L/\mathbb{Z}}^1 \oplus (O_L \otimes_{\mathbb{Z}} L^\times)) / \mathcal{N}, \quad (123)$$

where $\mathcal{N}$ is the O_L -submodule of the direct sum generated by elements of the form $(-da, a \otimes a)$ for $a \in O_L \setminus \{0\}$. For $a \in L^\times$, denote the element $(0, 1 \otimes a)$ by $d \log(a)$. For $r \geq 1$, let

$$\Omega_{O_L/\mathbb{Z}}^r(\log) = \bigwedge_{O_L}^r \Omega_{O_L/\mathbb{Z}}^1(\log), \quad (124)$$

and define

$$\Omega_{O_L}^r(\log) = \varprojlim \Omega_{O_L/\mathbb{Z}}^r(\log) / p^i \Omega_{O_L/\mathbb{Z}}^r. \quad (125)$$

Milnor K -theory then gives natural maps

$$d \log : K_r(L) \longrightarrow \Omega_{O_L}^r(\log) \quad (126)$$

which are defined by

$$\{a_1, \dots, a_r\} \longrightarrow d \log(a_1) \wedge \dots \wedge d \log(a_r). \quad (127)$$

For all $i \geq j \gg 0$, define the evaluation map

$$h_{i,j}^{(N)} : \mathcal{S}_K^{\dagger, N} \longrightarrow \mathcal{K}_j(\zeta_{p^i}) \quad (128)$$

determined by $h_{i,j}^{(N)}(\pi_K) = \pi_{K_i}$ and $h_{i,j}^{(N)}(T_k) = (T_k^{\frac{1}{p^j}})$ for all k . Note that $h_{j,j}^{(N)} = h_j^{(N)}$, which was defined above. Let

$$H_{i,j}^{(N)} : \widehat{K}_n(\mathcal{S}_K^{\dagger, N}) \longrightarrow \widehat{K}_n(\mathcal{K}_j(\zeta_{p^i})) \quad (129)$$

$$\mathcal{H}_{i,j}^{(N)} : \Omega_{\mathcal{S}_K^{\dagger, N}}^n(\log) \longrightarrow \Omega_{O_{\mathcal{K}_i}(\zeta_{p^i})}^n(\log) \quad (130)$$

be the (surjective) maps induced by the evaluation map $h_{i,j}^{(N)}$. If $i = j$, denote them by $H_i^{(N)}$ and $\mathcal{H}_i^{(N)}$.

Proposition 58. *We have a commutative diagram*

$$\begin{array}{ccc} \varprojlim \widehat{K}_n(\mathcal{K}_i) & \xrightarrow{d \log} & \varprojlim \Omega_{O_{\mathcal{K}_i}}^n(\log) \\ \downarrow T & \nearrow (\mathcal{H}_i^{(N)})_{i \geq 1} & \\ (\Omega_{\mathcal{S}_K^{\dagger, N}}^n(\log))^{\psi=1} & & \end{array}$$

Proof. Recall that we have a continuous homomorphism $g : \varprojlim \widehat{K}_n(\mathcal{K}_i) \rightarrow \widehat{K}_n(\mathbb{E}_{\mathcal{K}})$. Also, recall the following construction from the proof of Proposition 31:- For all $i \gg 0$, the evaluation map $h_i^{(N)}$ induces a homomorphism

$$H_i^{(N)} : \widehat{K}_n(\mathcal{T}_K^{\dagger, N}) \longrightarrow \widehat{K}_n(\mathcal{K}_i). \quad (131)$$

One can show that there exists $j_i^{(N)} \geq i - N - 1$ such that the image of $\ker(H_i^{(N)})$ in $\widehat{K}_n(\mathbb{E}_{\mathcal{K}})$ is the subgroup $\mathfrak{A}_i^{(N)}$ generated by the symbols of the form $\{a_1, \dots, a_n\}$ with $a_1 \in 1 + \pi_K^{j_i^{(N)}} \mathbb{E}_{\mathcal{K}}^+$, and we have commutative diagram

$$\begin{array}{ccccc} K_n(\mathbb{E}_{\mathcal{K}})/\mathfrak{A}_{i+1}^{(N)} & \longleftarrow & \widehat{K}_n(\mathcal{T}_K^{\dagger, N})/\ker(H_{i+1}^{(N)}) & \xrightarrow[\cong]{H_{i+1}^{(N)}} & \widehat{K}_n(\mathcal{K}_{i+1}) \\ \downarrow & & \downarrow N & & \downarrow N_{O_{\mathcal{K}_{m+1}}/O_{\mathcal{K}_m}} \\ K_n(\mathbb{E}_{\mathcal{K}})/\mathfrak{A}_i^{(N)} & \longleftarrow & \widehat{K}_n(\mathcal{T}_K^{\dagger, N})/\ker(H_i^{(N)}) & \xrightarrow[\cong]{H_i^{(N)}} & \widehat{K}_n(\mathcal{K}_i) \end{array}$$

Now $j_i^{(N)} \rightarrow +\infty$ as $i \rightarrow +\infty$, so we have $\varprojlim K_n(\mathbb{E}_{\mathcal{K}})/\mathfrak{A}_i^{(N)} = \widehat{K}_n(\mathbb{E}_{\mathcal{K}})$, and the above diagram induces a map

$$G : \varprojlim \widehat{K}_n(\mathcal{T}_K^{\dagger, N})/\ker(H_i^{(N)}) \longrightarrow \widehat{K}_n(\mathbb{E}_{\mathcal{K}}). \quad (132)$$

Now for all $i \gg 0$ we have a commutative diagram

$$\begin{array}{ccc} \mathcal{H}_{i+1}^{(n)} : \Omega_{\mathcal{S}_K^{\dagger, N}}^n(\log) & \longrightarrow & \Omega_{O_{\mathcal{K}_{i+1}}}^n(\log) \\ \downarrow \text{Tr}_{\phi}^{(n)} & & \downarrow \text{Tr} \\ \mathcal{H}_i^{(n)} : \Omega_{\mathcal{S}_K^{\dagger, N}}^n(\log) & \longrightarrow & \Omega_{O_{\mathcal{K}_i}}^n(\log) \end{array}$$

so $\text{Tr}_{\phi}^{(n)}(\ker(\mathcal{H}_{i+1}^{(N)})) \subset \ker(\mathcal{H}_i^{(N)})$ for all i , and we get a commutative diagram

$$\begin{array}{ccc} G : \varprojlim \widehat{K}_n(\mathcal{T}_K^{\dagger, N})/\ker(H_i^{(N)}) & \xrightarrow[\cong]{} & \varprojlim \widehat{K}_n(\mathcal{K}_i) \\ \downarrow d\log & & \downarrow d\log \\ (\mathcal{H}_i^{(n)})_{i \gg 0} : \varprojlim \Omega_{\mathcal{S}_K^{\dagger, N}}^n(\log)/\ker(\mathcal{H}_i^{(N)}) & \xrightarrow[\cong]{} & \varprojlim \Omega_{O_{\mathcal{K}_i}}^n(\log) \end{array}$$

Now let $x \in \varprojlim \widehat{K}_n(\mathcal{T}_K^{\dagger, N})/\ker(H_i^{(N)})$, and denote by $\bar{x}$ its image in $\widehat{K}_n(\mathbb{E}_{\mathcal{K}})$. Choose a sequence $(y^{(j)})$ of elements in $\tilde{K}_n(\mathcal{S}_K)$ such that for all $i \gg 0$ we have $\lambda(y^{(1)} \dots y^{(i)}) \cong \bar{x} \pmod{\mathfrak{A}_i^{(N)}}$. Note that we can choose $y^{(j)}$ such that

it can be written as a product of symbols with entries in $\mathcal{T}_K^{\dagger, N} \subset \mathcal{S}_K$. We will show that for all $i \gg 0$, $\psi^k(d \log(y^{(1)} \dots y^{(i+k)})) \rightarrow d \log(x) \pmod{\mathcal{H}_i^{(N)}}$ as $k \rightarrow +\infty$. Since $\mathcal{H}_i^{(N)}$ is the evaluation map on the differentials, this will prove the proposition. Now for each $j \gg 0$ there exist $y_1^{(j)}, y_2^{(j)} \in \widehat{K}_n(\mathcal{S}_K)$ such that $y_1^{(1)} \dots y_1^{(i)}$ is a lift of $x \pmod{\ker(H_i^{(N)})}$ and $y_2^{(1)} \dots y_2^{(i)} \in \ker(\lambda)$. By the definition of $y_1^{(j)}$, certainly $\psi^k d \log(y_1^{(1)} \dots y_1^{(i+k)}) = d \log(x) \pmod{\mathcal{H}_i^{(N)}}$. But $N^k(y_2^{(1)} \dots y_2^{(i+k)}) \rightarrow 1$ as $k \rightarrow +\infty$ by Lemma 23, so $\psi^k(d \log(y_2^{(1)} \dots y_2^{(i+k)})) \rightarrow 0$ as $k \rightarrow +\infty$, which finishes the proof. $\square$

7.2.2 The maps $\mathfrak{f}_{(\zeta_{p^i})_{i \geq 1}, m}$

Define

$$\lambda_{(\zeta_{p^i})_{i \geq 1}, m} : \varprojlim \widehat{K}_n(\mathcal{K}_m(\zeta_{p^i})) \xrightarrow{(\delta_{\mathcal{K}_m(\zeta_{p^i}), p^i}^{n-1})_{i \geq 1}} \varprojlim H^n(\mathcal{K}_m(\zeta_{p^i}), (\mathbb{Z}/p^i \mathbb{Z})(n)) \quad (133)$$

$$\xrightarrow{\zeta_{p^i}^{\otimes(-1)}} \varprojlim H^n(\mathcal{K}_m(\zeta_{p^i}), (\mathbb{Z}/p^i \mathbb{Z})(n-1)) \quad (134)$$

$$\xrightarrow{\text{trace}} \varprojlim H^n(\mathcal{K}_m, (\mathbb{Z}/p^i \mathbb{Z})(n-1)) \quad (135)$$

$$= H^n(\mathcal{K}_m, \mathbb{Z}_p(n-1)) \quad (136)$$

$$\longrightarrow H^n(\mathcal{K}_m, \mathbb{C}_p(n-1)) \cong \Omega_{\mathcal{K}_m}^{n-1}, \quad (137)$$

In [11], Kato defines a homomorphism

$$\mathfrak{f}_{(\zeta_{p^i})_{i \geq 1}, m} : \varprojlim \Omega_{\mathcal{K}_m(\zeta_{p^i})}^n(\log) \longrightarrow \Omega_{\mathcal{K}_m}^{n-1}, \quad (138)$$

and he shows the following result, which is crucial for our description of the dual exponential map:-

Theorem 59. *For $m \geq 1$, let $\mathfrak{d}_{(\zeta_{p^i})_{i \geq 1}, m} = \mathfrak{f}_{(\zeta_{p^i})_{i \geq 1}, m} \circ d \log$ be the map*

$$\varprojlim \widehat{K}_n(\mathcal{K}_m(\zeta_{p^i})) \xrightarrow{d \log} \varprojlim \Omega_{\mathcal{K}_m(\zeta_{p^i})}^n(\log) \quad (139)$$

$$\xrightarrow{\mathfrak{f}_{(\zeta_{p^i})_{i \geq 1}, m}} \Omega_{\mathcal{K}_m}^{n-1}. \quad (140)$$

Then

$$\lambda_{(\zeta_{p^i})_{i \geq 1}, m} = \mathfrak{d}_{(\zeta_{p^i})_{i \geq 1}, m}. \quad (141)$$

In the rest of this subsection, we will give a construction of the homomorphism $\mathfrak{f}_{(\zeta_{p^i})_{i \geq 1}, m}$, which we quote from Section 5 in [8]:-

Lemma 60. *For all $i \geq m$, we have*

$$p^{m-i} \text{Tr}_{\mathcal{K}_m(\zeta_{p^i})/\mathcal{K}_m}(O_{\mathcal{K}_m(\zeta_{p^i})}) \subset O_{\mathcal{K}_m} \quad (142)$$

where $\text{Tr}_{\mathcal{K}_m(\zeta_{p^i})/\mathcal{K}_m} : \mathcal{K}_m(\zeta_{p^i}) \rightarrow \mathcal{K}_m$ is the usual trace map.

Proof. Lemma 6.1.6 in [11]. $\square$

Lemma 61. *Let $i \geq m$, and define*

$$\kappa_i : O_{H_m(\zeta_{p^i})}/(p^i) \otimes_{O_K} \Omega_{O_K}^1 \longrightarrow \Omega_{O_{K_m(\zeta_{p^i})}}^2(\log) \quad (143)$$

$$x \otimes y \longrightarrow xy \wedge d \log(\zeta_{p^i}) \quad (144)$$

Then p^{e+m+2} kills $\ker(\kappa_i)$ and p^{e+m+1} kills $\text{coker}(\kappa_i)$.

Proof. Lemma 6.1.7 in [11]. $\square$

Now let $c = c_1 c_2 c_3$ for $c_1 = p^m$, $c_2 = p^{e+m+1}$ and $c_3 = p^{e+m+2}$, and define

$$f_{i,c} : \widehat{K}_n(K_m(\zeta_{p^i})) \longrightarrow O_{K_m}/p^i \otimes_{O_K} \Omega_{O_K}^{n-1} \quad (145)$$

as the composition

$$\widehat{K}_n(K_m(\zeta_{p^i})) \xrightarrow{d \log} \Omega_{K_m(\zeta_{p^i})}^n(\log) \quad (146)$$

$$\xrightarrow{c_2 \kappa_i^{-1} c_3} O_{K_m(\zeta_{p^i})}/p^i \otimes_{O_K} \Omega_{O_K}^{n-1} \quad (147)$$

$$\xrightarrow{c_1 p^{-i} \text{Tr}_{K_m(\zeta_{p^i})/K_m} \otimes \text{id}} O_{K_m}/p^i \otimes_{O_K} \Omega_{O_K}^{n-1} \quad (148)$$

Here, the map $c_2 \kappa_i^{-1} c_3$ is defined as follows:- For $x \in \Omega_{K_m(\zeta_{p^i})}^n(\log)$, let y be an element of $O_{K_m(\zeta_{p^i})}/p^i \otimes_{O_K} \Omega_{O_K}^{n-1}$ such that $c_3 x = \kappa_i(y)$. Then $c_2 \kappa_i^{-1} c_3(x) = c_2(y)$.

We can now define the map $\mathfrak{f}_{(\zeta_{p^i})_{i \geq 1}, m}$:-

Definition. Let

$$\mathfrak{f}_{(\zeta_{p^i})_{i \geq 1}, m} = c^{-1} \varprojlim_i f_{i,c}. \quad (149)$$

7.2.3 Lots of commutative diagrams

In view of Theorem 59, the proof of Theorem 57 reduces to showing the following result:-

Proposition 62. *The composition*

$$\varprojlim \widehat{K}_n(K_i) \xrightarrow{(N_{K_i/K_m(\zeta_{p^i})})_i} \varprojlim \widehat{K}_n(K_m(\zeta_{p^i})) \quad (150)$$

$$\xrightarrow{\mathfrak{f}_{(\zeta_{p^i})_{i \geq 1}, m}} \Omega_{K_m}^{n-1} \quad (151)$$

coincides with $\mathfrak{d}_m$.

Using the commutative diagram

$$\begin{array}{ccc}
\widehat{K}_n(\mathcal{K}_i) & \xrightarrow{d \log} & \Omega_{\mathcal{O}_{\mathcal{K}_i}}^n(\log) \\
\downarrow N_{\mathcal{K}_i/\mathcal{K}_m(\zeta_{p^i})} & & \downarrow \text{Tr}_{\mathcal{K}_i/\mathcal{K}_m(\zeta_{p^i})} \\
\widehat{K}_n(\mathcal{K}_m(\zeta_{p^i})) & \xrightarrow{d \log} & \Omega_{\mathcal{O}_{\mathcal{K}_m(\zeta_{p^i})}}^n(\log)
\end{array}$$

we see that to prove Proposition 62 it is sufficient to show the following result:-

Proposition 63. *The map*

$$\varprojlim \widehat{K}_n(\mathcal{K}_i) \xrightarrow{d \log} \varprojlim \Omega_{\mathcal{O}_{\mathcal{K}_i}}^n(\log) \quad (152)$$

$$\xrightarrow{\text{Tr}_{\mathcal{K}_i/\mathcal{K}_m(\zeta_{p^i})}} \varprojlim \Omega_{\mathcal{O}_{\mathcal{K}_m(\zeta_{p^i})}}^n(\log) \quad (153)$$

$$\xrightarrow{\mathfrak{f}_{(\zeta_{p^i})_{i \geq 1}, m}} \Omega_{\mathcal{K}_m}^{n-1} \quad (154)$$

agrees with $\mathfrak{d}_m$.

Proof. Define homomorphisms $\sigma, \tau : \mathcal{S}_K^{\dagger, N} \rightarrow \mathcal{S}_K^{\dagger, n+1}$, which are determined by

$$\sigma : \pi_K \longrightarrow \phi(\pi_K), \quad (155)$$

$$\tau : T_i \longrightarrow T_i^p \text{ for all } i. \quad (156)$$

Note that $\phi = \sigma \circ \tau$. As before, these homomorphisms then induce trace maps

$$\text{Tr}_{\sigma}^{(n)}, \text{Tr}_{\tau}^{(n)} : \Omega_{\mathcal{S}_K^{\dagger, N}}^n(\log) \longrightarrow \Omega_{\mathcal{S}_K^{\dagger, n}}^n(\log). \quad (157)$$

Explicit computation shows that

$$\begin{aligned}
\text{Tr}_{\sigma}^{(n)}\left(\frac{a}{\pi_K} \bigwedge d \log(T_i) \wedge d \log(\pi_K + 1)\right) &= \frac{1}{p} \frac{\text{Tr}_{\sigma}(p \cdot a \omega)}{\pi_K} \bigwedge d \log(T_i) \wedge d \log(\pi_K + 1) \\
\text{Tr}_{\tau}^{(n)}\left(\frac{b}{\pi_K} \bigwedge d \log(T_i) \wedge d \log(\pi_K + 1)\right) &= \frac{1}{p^{n-1}} \frac{\text{Tr}_{\tau}(p \cdot b \omega)}{\pi_K} \bigwedge d \log(T_i) \wedge d \log(\pi_K + 1)
\end{aligned}$$

where

$$\omega = \frac{\phi(\pi_K)}{\pi_K} \frac{\phi(\pi_K) + 1}{\pi_K + 1} \left(\frac{d}{d\pi_K} \phi(\pi_K + 1) \right)^{-1}, \quad (158)$$

and we have a commutative diagram

$$\begin{array}{ccc}
\Omega_{\mathcal{S}_K^{\dagger, N}}^n(\log) & \xrightarrow{(\text{Tr}_{\sigma}^{(n)})^{i-m}} & \Omega_{\mathcal{S}_K^{\dagger, n}}^n(\log) \\
\downarrow \mathcal{H}_{i, m}^{(N)} & & \downarrow \mathcal{H}_m^{(N)} \\
\Omega_{\mathcal{O}_{\mathcal{K}_m(\zeta_{p^i})}}^n(\log) & \xrightarrow{\text{Tr}_{\mathcal{K}_i/\mathcal{K}_m(\zeta_{p^i})}^{(n)}} & \Omega_{\mathcal{O}_{\mathcal{K}_i}}^n(\log)
\end{array}$$

Also, τ induces a trace map

$$\mathrm{Tr}_\tau : \frac{\mathcal{S}_K^{\dagger,N}}{\pi_K} \longrightarrow \frac{\mathcal{S}_K^{\dagger,N}}{\pi_K},$$

and we have a commutative diagram

$$\begin{array}{ccc} \frac{\mathcal{S}_K^{\dagger,N}}{\pi_K} & \xrightarrow{(\mathrm{Tr}_\tau)^{i-m}} & \frac{\mathcal{S}_K^{\dagger,N}}{\pi_K} \\ \downarrow h_{i,m}^{(N)} & & \downarrow h_i^{(N)} \\ \mathcal{K}_m(\zeta_{p^i}) & \xrightarrow{\mathrm{Tr}_{\mathcal{K}_m(\zeta_{p^i})/\mathcal{K}_m}} & \mathcal{K}_m \end{array}$$

Here, $\mathrm{Tr}_\tau(\frac{c}{\pi_K}) = \frac{\mathrm{Tr}_\tau(\omega \cdot c)}{\pi_K}$, where $\mathrm{Tr}_\tau : \mathcal{S}_K^{\dagger,N} \rightarrow \mathcal{S}_K^{\dagger,N}$ is the usual trace map.

Recalling the definition of $\mathfrak{f}_{(\zeta_{p^i})_{i \geq 1}, m}$ and combining the above commutative diagrams with Proposition 58 shows that the map given by equations (150) and (144) is also given by the composition

$$\varprojlim \widehat{K}_n(\mathcal{K}_i) \longrightarrow (\Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log))^{\psi=1} \quad (159)$$

$$\hookrightarrow \Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log) \xrightarrow{(\mathrm{Tr}_\sigma^{(n)})^{i-m}} \Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log) \quad (160)$$

$$\xrightarrow{\mathfrak{pr}} \frac{\mathcal{S}_K^{\dagger,N}}{\pi_K} \bigwedge d\log(T_i) \quad (161)$$

$$\xrightarrow{(\mathrm{Tr}_\tau)^{i-m}} \frac{\mathcal{S}_K^{\dagger,N}}{\pi_K} \bigwedge d\log(T_i) \quad (162)$$

$$\longrightarrow \Omega_{\mathcal{K}_m}^{n-1} \quad (163)$$

where the map $\mathfrak{pr}$ is given by

$$\mathfrak{pr} : f \bigwedge d\log(T_i) \wedge d\log(\pi_K + 1) \longrightarrow f \bigwedge d\log(T_i)$$

and the last arrow is defined as

$$f \bigwedge d\log(T_i) \longrightarrow \frac{1}{p^m} h_m^{(N)}(f) \bigwedge d\log(h_m^{(N)}(T_i)). \quad (164)$$

But $\mathrm{Tr}_\tau^{(n)} \circ \mathrm{Tr}_\sigma^{(n)} = \mathrm{Tr}_\phi^{(n)}$ and $\mathrm{Tr}_\tau(\mathfrak{pr}(x)) = \mathfrak{pr}(\mathrm{Tr}_\tau^{(n)}(x))$ for all $x \in \Omega_{\mathcal{S}_K^{\dagger,N}}^n(\log)$, which shows that this composition also gives the map $\mathfrak{d}_m$. $\square$

References

- [1] L. Berger, *Représentations p-adiques et équations différentielles*, Invent. Math. **148** (2002), 219–284.

- [2] F. Cherbonnier and P. Colmez, *Représentations p -adiques surconvergentes*, Invent. Math. **133** (1998), no. 3, 581–611.
- [3] F. Cherbonnier and P. Colmez, *Théorie d’Iwasawa des représentations p -adiques d’un corps local*, J. Amer. Math. Soc. **12** (1999), no. 1, 241–268.
- [4] J. Coates and A. Wiles, *On p -adic L -functions and Elliptic Units*, J. Austral. Math. Soc. (Series A) (1978), no. 26, 1–25.
- [5] R. Coleman, *Division values in local fields*, Invent. Math. (1979), no. 53, 91–116.
- [6] P. Colmez, *p -adic L -functions and Fontaine’s rings*, Course given at Tsinghua, 2005.
- [7] T. Fukaya, *Explicit reciprocity laws for p -divisible groups over higher dimensional local fields*, Journal für die reine und ang. Math. (2001), no. 531, 61–119.
- [8] T. Fukaya, *The theory of Coleman power series for K_2* , Journal of Alg. Geometry (2003), no. 12, 1–80.
- [9] K. Kato, *A generalisation of local class field theory by using K -groups I*, J. Fac. Sci. Univ. of Tokyo, Sec. IA (1979), no. 26, 303–376.
- [10] K. Kato, *A generalisation of local class field theory by using K -groups II*, J. Fac. Sci. Univ. of Tokyo, Sec. IA (1980), no. 27, 603–683.
- [11] K. Kato, *Generalized explicit reciprocity laws*, Advanced Studies in Contemporary Mathematics 1 (1999), 57–126.
- [12] M. Kurihara, *The exponential homomorphism for Milnor K -groups and explicit reciprocity law*, Journal fuer reine und ang. Math. (1998), no. 498, 201–221.
- [13] A.J. Scholl, *Higher fields of norms*, (2005), preprint.
- [14] C. Weibel, *An introduction to algebraic K -theory*, (2002), online lecture notes.